M
8918

22500288168

R. A. Young.

June 23d 1899.



Syrup. Ipt.

Hydrocodèine

103 Tamponnets

PHARMACOPŒIA

OF

THE MIDDLESEX HOSPITAL.

LONDON:
MITCHELL AND HUGHES,
140 WARDOUR STREET, W.
1899.

24 880 257

95400 / 29266

WELLCOME INSTITUTE LIBRARY

Coll. We[illegible]

Coll. Gen. Coll.

No. QV 740

M

8918

WMC P

TABLE OF CONTENTS.

	Pages
Pharmacopœia :—	
A. General Formulæ	1 to 35
B. Formulæ for Children ..	36 „ 41
C. Formulæ for the Eye Department	42 „ 44
D. Formulæ for the Ear and Throat Department ..	45 „ 48
E. Formulæ for the Skin Department	49 „ 62
Treatment of Poisoning	63 „ 70
Table of Active Ingredients	71 „ 77
Preparations not included in the British Pharmacopœia	78 „ 79
Table for Preparing Effervescing Draughts	80
Posological Table	81
Table of Doses	81 „ 97
Instructions for Case-Taking in Medical Wards	101 „ 111
Instructions for Case-Taking in Surgical Wards	115 „ 124
Diet Table	125

PHARMACOPŒIA

OF

THE MIDDLESEX HOSPITAL.

A.

GENERAL FORMULÆ.

NOTE.—Preparations not included in the "British Pharmacopœia" are indicated in *italics*, and their method of preparation, when not elsewhere given, will be found on p. 78.

CHARTA POTASSII NITRATIS.

℞ Potassii Nitratis .. .. oz. 1.
Aquæ destillatæ.. .. .. fl. oz. iv.

Dissolve. Soak pieces of thin bibulous paper 8 inches square in this solution and dry.

One to be burnt for each inhalation.

General Formulæ.

CONFECTIO SENNÆ.

℞ Pulveris Sennæ.. .. .. oz. ½.
Pulveris Coriandri
Pulveris Jalapæ
Pulveris Potassæ Tartratis Acidæ
Pulveris Zingiberis
Pulveris Glycyrrhizæ, āā .. gr. 60.
Theriacæ .. .. .. oz. 3.

Mix. Dose, gr. 60 to gr. 120.

CONFECTIO SULPHURIS.

℞ Sulphuris Sublimati
Potassii Tartratis Acidæ, āā .. oz. ½.
Confectionis Sennæ .. .. oz. 2.
Theriacæ .. .. .. q. s.

Mix. Dose, gr. 60 to gr. 120.

EMULSIO OLEI MORRHUÆ.

℞ Olei Morrhuæ .. .. .. fl. oz. x.
Pulveris Tragacanthæ.. .. gr. 50.
Spiritûs Chloroformi .. .. fl. dr. iij.
Glycerini .. .. .. fl. dr. iv.
Olei Amygdalæ Essentii .. min. x.
Aquam destillatam .. ad O j.

Mix. Make an Emulsion. Dose, oz. fl. ss. to oz. fl. j.

General Formulæ.

ENEMA NUTRIENS (A).

℞ Peptonized Beef (Pepsin Peptone) .. .. gr. 60.
Extract of Malt .. .. .. gr. 60.
Gruel .. .. fl. oz. ij.

ENEMA NUTRIENS (B).

℞ Peptonized Beef (Pepsin Peptone) .. .. gr. 60.
Warm Water (98° Fht.) .. fl. oz. ij.

ENEMA OLEI OLIVÆ.

℞ Olei Olivæ .. .. .. fl. oz. iv.
Enema Simplex .. .. ad O j.

Mix.

ENEMA OLEI RICINI.

℞ Olei Ricini .. .. .. fl. oz. j.
Enema Simplex .. .. ad O j.

Mix.

General Formulæ.

ENEMA RUTÆ.

℞ Olei Rutæ .. .. .. min xx.
Mucilaginis Acaciæ .. .. fl. dr. ij.
Enema Simplex .. .. ad fl. oz. vj.

Mix.

ENEMA SIMPLEX.

℞ Saponis Mollis .. .. .. oz. 1.
Aquæ ferventis .. .. .. O j.

Dissolve.

ENEMA TEREBINTHINÆ.

℞ Olei Terebinthinæ .. .. fl. oz. ss.
Olei Ricini .. .. .. fl. oz. j.
Decocti Hordei vel Mucilaginis Amyli .. .. O j.

Mix.

FOTUS BELLADONNÆ.

℞ Extracti Belladonnæ Viride .. gr. 60.
Aquæ destillatæ .. .. .. O j.

Dissolve and filter.
To be made hot and used as a fomentation.

General Formulæ.

FOTUS PAPAVERIS.

℞ Papaveris Capsulæ .. .. oz. j.
Aquæ destillatæ .. .. .. O j.

Boil for a quarter of an hour in a covered vessel, strain, and add water to make up to a pint.

FUMIGATIO CHLORI.

℞ Sodii Chloridi .. .. .. oz. j.
Manganesii Oxidi Nigri.. .. oz. ¾.

Mix. Chlorine is evolved on the addition of Sulphuric Acid.

To be used only for disinfecting purposes.

GLYCERINUM BELLADONNÆ.

℞ Extracti Belladonnæ Viride .. oz. ½.
Aquæ destillatæ .. .. .. fl. dr. j.
Glycerini .. .. fl. dr. iij.

Rub the extract with the water, then add the glycerine and mix well together.

GLYCERINUM FERRI PERCHLORIDI.

℞ Liquoris Ferri Perchloridi
Glycerini, āā .. .. .. fl. oz. j.

Mix.

General Formulæ.

HAUSTUS ACIDI GALLICI.

℞ Acidi Gallici .. .. .. gr. 10.
Acidi Sulphurici Aromatici .. min. xv.
Spiritûs Chloroformi .. .. min. x.
Aquam destillatam .. ad fl. oz. j.
Mix.

HAUSTUS ÆTHERIS COMPOSITUS.

℞ Ammonii Carbonatis .. .. gr. 3.
Spiritûs Ætheris.. .. .. fl. dr. ss.
Spiritûs Chloroformi .. .. min. x.
Aquam destillatam .. ad fl. oz. j.
Mix.

HAUSTUS AMMONII ACETATIS.

℞ Liquoris Ammonii Acetatis .. fl. dr. iij.
Aquam destillatam .. ad fl. oz. j.
Mix.

HAUSTUS AMMONII ACETATIS CUM FERRO.

℞ Liquoris Ammonii Acetatis .. fl. dr. ij. ss.
Liquoris Ferri Perchloridi .. min. x.
Acidi Acetici diluti .. .. fl. dr. ss.
Aquam destillatam .. ad fl. oz. j.
Mix.

General Formulæ.

HAUSTUS APOMORPHINÆ COMPOSITUS.

℞	Apomorphinæ Hydrochloridi	.. gr. $\frac{1}{10}$.
	Acidi Hydrochlorici diluti	.. min. j.
	Syrupi Scillæ	.. fl. dr. j.
	Olei Terebinthinæ ..	.. min. x.
	Mucilaginis Acaciæ ..	.. q. s.
	Spiritûs Ætheris.. ..	.. min. x.
	Aquam destillatam ..	ad fl. oz. j.

Mix.

HAUSTUS BISMUTHI.

℞	Bismuthi Carbonatis	
	Magnesiæ Carbonatis Levis, āā..	gr. 10.
	Tincturæ Calumbæ ..	.. min. xx.
	Aquam destillatam ..	ad fl. oz. j.

Mix.

HAUSTUS CALCIS HYPOPHOSPHITIS COMPOSITUS.

℞	Calcis Hypophosphitis ..	.. gr. 5.
	Syrupi Ferri Phosphatis	.. fl. dr. j.
	Glycerini..	.. fl. dr. j.
	Aquam destillatam ..	ad fl. oz. j.

Mix.

General Formulæ.

HAUSTUS CARYOPHYLLI COMPOSITUS.

℞ Sodii Bicarbonatis .. .. gr. 10.
Spiritûs Ammoniæ Aromatici .. min. xx.
Spiritûs Chloroformi .. .. min. x.
Infusum Caryophylli .. ad fl. oz. j.
Mix.

HAUSTUS CHLOROFORMI COMPOSITUS.

℞ Spiritûs Chloroformi .. .. min. xv.
Tincturæ Camphoræ Compositæ min. xx.
Syrupi Scillæ .. .. .. fl. dr. ss.
Aquam destillatam .. ad fl. oz. j.
Mix.

HAUSTUS CINCHONÆ ACIDUS.

℞ Acidi Nitro-hydrochlorici diluti min. x.
Spiritûs Chloroformi .. .. min. v.
Decoctum Cinchonæ .. ad fl. oz. j.
Mix.

HAUSTUS CINCHONÆ CUM AMMONIÂ.

℞ Ammonii Carbonatis .. .. gr. 5.
Spiritûs Chloroformi .. .. min. v.
Decoctum Cinchonæ .. ad fl. oz. j.
Mix.

General Formulæ.

HAUSTUS COPAIBÆ.

℞	Copaibæ	.. fl. dr. ss.
	Liquoris Potassæ ..	.. min. xx.
	Mucilaginis Acaciæ ..	.. fl. dr. ij.
	Aquam Carui	ad fl. oz. j.

Mix.

HAUSTUS COPAIBÆ RESINÆ.

℞	*Tincturæ Copaibæ Resinæ	.. fl. dr. ss.
	Spiritûs Chloroformi ..	.. min. x.
	Syrupi Zingiberis	
	Mucilaginis Acaciæ, āā..	.. fl. dr. j.
	Aquam destillatam ..	ad fl. oz. j.

Mix.

* This tincture is prepared by liquefying 480 grains of the resin by the heat of a water bath, and mixing with it sufficient rectified spirit of wine, previously made warm, to measure 3 fluid ounces when the *temperature falls to* 60° F. Three minims contain one grain of the resin.

HAUSTUS CRETÆ COMPOSITUS.

℞	Cretæ Preparatæ ..	.. gr. 15.
	Mucilaginis Acaciæ ..	.. fl. dr. j.
	Tincturæ Catechu ..	.. min. xx.
	Aquam destillatam ..	ad fl. oz. j.

Mix.

General Formulæ.

HAUSTUS EMETICUS PURGANS.

℞	Antimonii Tartarati ..	.. gr. $\frac{1}{6}$. $\frac{3}{4}$
	Magnesii Sulphatis ..	.. gr. 60.
	Aquam destillatam ..	ad fl. oz. j.

Mix.

HAUSTUS ERGOTÆ.

℞	Extracti Ergotæ Liquidi	.. fl. dr. ss.
	Spiritûs Chloroformi ..	.. min. v.
	Aquam destillatam ..	ad fl. oz. j.

Mix.

HAUSTUS FERRI ET AMMONII CITRATIS.

℞	Ferri et Ammonii Citratis	.. gr. 5.
	Spiritûs Chloroformi ..	.. min. v.
	Infusum Quassiæ ..	ad fl. oz. j.

Mix.

HAUSTUS FERRI CUM MAGNESIÂ.

℞	Ferri Sulphatis	.. gr. 1.
	Magnesii Sulphatis ..	.. gr. 20.
	Acidi Sulphurici diluti ..	.. min. x.
	Aquam Menthæ Piperitæ	ad fl. oz. j.

Mix.

General Formulæ.

HAUSTUS FERRI PERCHLORIDI.

℞ Liquoris Ferri Perchloridi .. min. xv.
Spiritûs Chloroformi .. .. min. x.
Glycerini .. .. min. xx.
Aquam destillatam .. ad fl. oz. j.
Mix.

HAUSTUS FILICIS MARIS.

℞ Extracti Filicis Liquidi .. .. fl. dr. j.
Pulveris Tragacanthæ Compositi gr. 40.
Aquam Menthæ Piperitæ ad fl. oz. j.
Mix.

HAUSTUS GENTIANÆ ACIDUS.

℞ Acidi Nitro-hydrochlorici diluti min. x.
Spiritûs Chloroformi .. .. min. v.
Infusum Gentianæ .. ad fl. oz. j.
Mix.

HAUSTUS GENTIANÆ ALKALINUS.

℞ Potassii Bicarbonatis .. .. gr. 10.
Tincturæ Zingiberis .. .. min. x.
Spiritûs Chloroformi .. .. min. v.
Infusi Rhei .. .. .. fl. dr. ss.
Infusum Gentianæ .. ad fl. oz. j.
Mix.

General Formulæ.

HAUSTUS HYDRARGYRI IODIDI.

℞ Potassii Iodidi .. .. .. gr. 5.
Liquoris Hydrargyri Perchloridi fl. dr. j.
Spiritûs Chloroformi .. .. min. v.
Infusum Chiratæ .. ad fl. oz. j.

Mix.

HAUSTUS IPECACUANHÆ COMPOSITUS.

℞ Vini Ipecacuanhæ .. .. min. x.
Tincturæ Camphoræ Compositæ fl. dr. ss.
Liquoris Ammonii Acetatis .. fl. dr. iij.
Aquam destillatam .. ad fl. oz. j.

Mix.

HAUSTUS IPECACUANHÆ CUM AMMONIÂ.

℞ Ammonii Carbonatis .. .. gr. 4.
Vini Ipecacuanhæ
Spiritûs Chloroformi, āā .. min. x.
Syrupi Rhœados.. .. .. fl. dr. ss.
Aquam destillatam .. ad fl. oz. j.

Mix.

General Formulæ.

HAUSTUS IPECACUANHÆ CUM SCILLÂ.

℞ Vini Ipecacuanhæ .. .. min. x.
Aceti Scillæ .. .. .. fl. dr. ss.
Glycerini .. .. min. xxx.
Aquam Menthæ Viridis.. ad fl. oz. j.
Mix.

HAUSTUS MAGNESIÆ COMPOSITUS.

℞ Magnesii Carbonatis Levis .. gr. 10.
Magnesii Sulphatis .. .. gr. 60.
Aquæ Menthæ Piperitæ.. .. fl. oz. j.
Mix.

HAUSTUS OLEI RICINI.

℞ Olei Ricini .. .. .. fl. oz. ss.
Liquoris Potassæ .. .. min. xv.
Aquam Menthæ Piperitæ ad fl. oz. j.
Mix.

HAUSTUS POTASSII CHLORATIS CUM FERRO.

℞ Potassii Chloratis .. .. gr. 10.
Liquoris Ferri Perchloridi .. min. x.
Acidi Hydrochlorici diluti .. min. v.
Glycerini .. .. min. xxx.
Aquam destillatam .. ad fl. oz. j.
Mix.

General Formulæ.

HAUSTUS POTASSII CITRATIS.

℞ Potassii Citratis .. .. .. gr. 30.
Tincturæ Aurantii .. .. min. v.
Aquam destillatam .. ad fl. oz. j.

Mix.

HAUSTUS POTASSII CITRATIS EFFERVESCENS.

℞ Potassii Bicarbonatis .. .. gr. 20.
Aquæ destillatæ .. .. .. fl. oz. ss.

Dissolve.

℞ Acidi Citrici .. .. .. gr. 14.
Aquæ destillatæ .. .. .. fl. oz. ss.

Dissolve.

Mix the two solutions. To be taken while effervescing.

HAUSTUS POTASSII IODIDI.

℞ Potassii Iodidi .. .. .. gr. 5.
Spiritûs Ammoniæ Aromatici .. min. xx.
Decoctum Cinchonæ .. ad fl. oz. j.

Mix.

General Formulæ.

HAUSTUS POTASSII IODIDI ALKALINUS.

℞ Potassii Iodidi .. .. .. gr. 2.
Potassii Bicarbonatis .. .. gr. 15.
Spiritûs Chloroformi .. .. min. v.
Infusum Gentianæ .. ad fl. oz. j.

Mix.

HAUSTUS POTASSII IODIDI CUM COLCHICO.

℞ Potassii Iodidi .. .. .. gr. 5.
Potassii Bicarbonatis .. .. gr. 10.
Vini Colchici .. .. .. min. x.
Spiritûs Chloroformi .. .. min. v.
Infusum Cascarillæ .. ad fl. oz. j.

Mix.

HAUSTUS QUININÆ.

℞ Quininæ Sulphatis .. .. gr. 1.
Acidi Sulphurici diluti .. .. min. j.
Aquam destillatam .. ad fl. oz. j.

Mix.

General Formulæ.

HAUSTUS QUININÆ AMMONIATUS.

℞ Tincturæ Quininæ Ammoniatæ fl. dr. j.
Spiritûs Chloroformi .. .. min. x.
Aquam destillatam .. ad fl. oz. j.
Mix.

HAUSTUS QUININÆ CUM FERRO.

℞ Quininæ Sulphatis
Ferri Sulphatis, āā .. .. gr. 1.
Acidi Sulphurici diluti .. .. min. ij.
Glycerini .. .. min. xx.
Aquam destillatam .. ad fl. oz. j.
Mix.

HAUSTUS SALINUS.

℞ Potassii Nitratis.. .. .. gr. 10.
Liquoris Ammonii Acetatis .. fl. dr. ij. ss.
Aquam Camphoræ .. ad fl. oz. j.
Mix.

HAUSTUS SCILLÆ ACIDUS.

℞ Aceti Scillæ .. .. .. min. xv.
Spiritûs Ætheris Nitrosi .. fl. dr. ss.
Liquoris Ammonii Acetatis .. fl. dr. ij.
Aquam Carui .. .. ad fl. oz. j.
Mix.

General Formulæ.

HAUSTUS SCOPARII.

℞ Potassii Acetatis.. .. .. gr. 20.
Tincturæ Scillæ .. .. .. min. x.
Spiritûs Ætheris Nitrosi .. fl. dr. ss.
Infusum Scoparii .. ad fl. oz. j.

Mix.

HAUSTUS SENEGÆ COMPOSITUS.

℞ Ammonii Carbonatis .. .. gr. 3.
Tincturæ Scillæ .. .. .. min. x.
Olei Pimentæ .. .. .. min. j.
Spiritûs Chloroformi .. .. min. x.
Infusum Senegæ.. .. ad fl. oz. j.

Mix.

HAUSTUS SENNÆ COMPOSITUS.

℞ Magnesii Sulphatis .. .. gr. 120.
Sodii Sulphatis .. .. .. gr. 30.
Extracti Glycyrrhizæ .. .. gr. 10.
Olei Pimentæ .. .. .. min. j.
Essentiæ Menthæ Piperitæ .. min. v.
Infusum Sennæ .. .. ad fl. oz. j.

Mix.

General Formulæ.

HAUSTUS SODII SALICYLATIS.

℞ Sodii Salicylatis .. .. .. gr. 15.
Olei Limonis .. .. .. min. j.
Spiritûs Chloroformi .. .. min. iv.
Aquam destillatam .. ad fl. oz. j.

Mix.

HAUSTUS STRYCHNINÆ COMPOSITUS.

℞ Quininæ Sulphatis
Zinci Sulphatis, āā .. .. gr. 1.
Liquoris Strychninæ Hydrochloridi .. .. .. min. iv.
Acidi Sulphurici diluti .. .. min. ij.
Aquam destillatam .. ad fl. oz. j.

Mix.

HAUSTUS STRYCHNINÆ CUM MAGNESIÂ.

℞ Ferri Sulphatis .. .. .. gr. 2.
Magnesii Sulphatis .. .. gr. 15.
Liquoris Strychninæ Hydrochloridi .. .. .. min. iv.
Spiritûs Chloroformi .. .. min. x.
Aquam Menthæ Piperitæ ad fl. oz. j.

Mix.

For Chr. Tab. Nephritis

Ferrum Tart. gr x
Acid Bor. Sanitate gr x
Sp. Mitis Mit. m xx.
ad 3j.

Heat Stych. ē Lemin

Lem. Sulph. gtt i
Acid Nitroby. dil. m x
Liq. Stych. hydrochlor. m iii
Syrupi Aurantii 3j
Aq. Chloroformi ad 3j.

General Formulæ.

HAUSTUS VALERIANÆ COMPOSITUS.

℞ Tincturæ Valerianæ Ammoniatæ min. xx.
Spiritûs Ammoniæ Fœtidi .. min. xv.
Infusum Quassiæ .. ad fl. oz. j.

Mix.

INJECTIO ALUMINIS.

℞ Aluminis .. .. gr. 5.
Aquæ destillatæ .. .. .. fl. oz. j.

Dissolve.

INJECTIO CUPRI SULPHATIS.

℞ Cupri Sulphatis .. .. .. gr. 1.
Aquæ destillatæ .. .. .. fl. oz. j.

Dissolve.

INJECTIO MORPHINÆ HYPODERMICA FORTIOR.

℞ Morphinæ Acetatis .. .. gr. 80.
Aquæ destillatæ .. .. .. fl. oz. j.

Rub the Acetate gradually with the water, and add a few drops of Acetic Acid, if necessary for perfect solution. Min. vj contain gr. 1 of Acetate of Morphine.

General Formulæ.

INJECTIO MORPHINÆ ET ATROPINÆ HYPODERMICA.

℞ Atropinæ Sulphatis .. .. gr. 1.
Injectionis Morphinæ Hypodermicæ .. .. fl. dr. x.

Dissolve.

Dose, two to eight minims. Six minims contain $\frac{3}{10}$ths of a grain of Tartrate of Morphine and $\frac{1}{100}$th of a grain of Sulphate of Atropine.

INJECTIO PLUMBI.

℞ Liquoris Plumbi Subacetatis .. fl. dr. j.
Aquam destillatam .. ad O j.

Mix.

INJECTIO QUERCÛS COMPOSITA.

℞ Aluminis
Zinci Sulphatis
Acidi Borici, āā .. .. .. gr. 80.
Decocti Quercûs
Aquæ destillatæ, āā .. .. O ss.

Mix.

General Formulæ.

INJECTIO ZINCI CHLORIDI.

℞	Zinci Chloridi	.. gr. 1.
	Acidi Hydrochlorici diluti	.. min. j.
	Aquam destillatam ..	ad fl. oz. j.

Mix.

INJECTIO ZINCI PERMANGANATIS.

℞	Zinci Permanganatis ..	.. gr. 1.
	Aquæ destillatæ	.. fl. oz. viij.

Dissolve.

INJECTIO ZINCI SULPHATIS.

℞	Zinci Sulphatis	.. gr. 2.
	Aquæ destillatæ	.. fl. oz. j.

Dissolve.

LINCTUS.

℞	Syrupi Rhœados.. ..	.. min. xx.
	Syrupi Scillæ	.. min. xv.
	Succi Limonis	.. min. xv.
	Syrupum	ad fl. dr. j.

Mix. Dose, fl. dr. ss. to fl. dr. j.
To be slowly sucked from a tea-spoon.

General Formulæ.

LINCTUS MORPHINÆ CUM CHLOROFORMO.

℞ Liquoris Morphinæ Acetatis .. min. viij.
Chloroformi .. .. .. min. ⅓.
Succi Limonis .. .. .. min. xv.
Glycerinum .. .. ad fl. dr. j.

Mix. Dose, fl. dr. ss. to fl. dr. j.
To be slowly sucked from a tea-spoon.

LINIMENTUM CALCIS (Carron Oil).

℞ Olei Lini
Liquoris Calcis, āā .. .. fl. oz. iv.

Mix.

LINIMENTUM SAPONIS MOLLIS.

℞ Saponis Mollis .. .. .. oz. ss.
Aquæ ferventis .. .. .. fl. oz. v.

Mix.

LINIMENTUM SAPONIS CUM OPIO.

℞ Tincturæ Opii .. .. .. fl. dr. ij.
Linimenti Saponis Mollis .. fl. dr. vj.

Mix.

General Formulæ.

LINIMENTUM TEREBINTHINÆ CUM AMMONIÂ.

℞ Linimenti Terebinthinæ
Liquoris Ammoniæ, āā.. .. fl. oz. j. ss.
Olei Cajuputi .. .. .. fl. dr. ss.
Olei Olivæ .. .. .. fl. oz. iv.

Mix.

LOTIO ACIDI HYDROCYANICI COMPOSITA.

℞ Boracis .. .. gr. 40.
Acidi Hydrocyanici diluti .. fl. dr. ss.
Aquam destillatam .. ad fl. oz. x.

Mix.

LOTIO ACIDI NITRICI CUM OPIO.

℞ Acidi Nitrici diluti .. .. fl. dr. j.
Extracti Opii Liquidi .. .. fl. dr. iij.
Aquam destillatam .. ad fl. oz. viij.

Mix.

LOTIO EVAPORANS.

℞ Spiritûs Rectificati .. .. fl. oz. ij.
Olei Lavandulæ .. .. .. min. xx.
Aquam destillatam .. ad O j.

Mix.

General Formulæ.

LOTIO FERRI SULPHATIS.

℞ Ferri Sulphatis .. .. .. gr. 100.
Acidi Sulphurici diluti .. .. min. c.
Aquam destillatam .. ad O j.
Mix.

LOTIO NIGRA.

℞ Hydrargyri Subchloridi .. gr. 60.
Mucilaginis Tragacanthæ .. fl. oz. ½.
Liquorem Calcis.. .. ad O j.
Mix.

LOTIO PLUMBI CUM OPIO.

℞ Extracti Opii Liquidi .. .. fl. oz. ss.
Liquorem Plumbi Subacetatis dilutem .. .. ad O j.
Mix.

LOTIO RUBRA.

℞ Zinci Sulphatis .. .. .. gr. 1.
Spiritûs Rosmarini
Tincturæ Lavandulæ Compositæ,
āā .. .. min. xv.
Aquam destillatam .. ad fl. oz. j.
Mix.

General Formulæ.

LOTIO ZINCI CHLORIDI.

℞ Zinci Chloridi .. .. .. gr. 20.
Acidi Hydrochlorici diluti .. min. xx.
Aquam destillatam .. ad O j.

Mix.

PASTA AMYLI IODIDI.

℞ Pulveris Amyli .. .. .. oz. j.
Glycerini .. .. fl. oz. ij.
Aquæ destillatæ .. .. .. fl. oz. vj.

Boil together, and when nearly cold add half a fluid ounce of Liquor Iodi Fortis (P.B.).

PASTA POTASSÆ CUM CALCE.

(Vienna Paste.)

℞ Potassæ Causticæ
Calcis Ustæ Recentis, āā, partes æquales.

Mix. To be made into a paste with a little Rectified Spirit.

General Formulæ.

PASTA ZINCI CHLORIDI CUM OPIO.

℞	Zinci Chloridi	.. oz. 16.
	Pulveris Opii	.. oz. 1½.
	Acidi Hydrochlorici ..	.. fl. dr. vj.
	Farinæ Tritici	.. gr. 60.
	Aquam bullientem ..	ad O j.

Macerate the Opium in twelve ounces of the boiling water for twelve hours, add the Acid and filter, then dissolve the Chloride of Zinc in the filtered liquid and make up to twenty ounces with distilled water. Add the flour. Mix smoothly in a mortar and heat over a water-bath until of a proper consistence.

PIGMENTUM IODI.

℞	Liquoris Iodi Fortis	
	Tincturæ Iodi, āā ..	.. oz. ss.

PILULA ALOES CUM NUCE VOMICÂ.

℞	Extracti Nucis Vomicæ..	.. gr. ¼.
	Extracti Aloës Barbadensis	.. gr. 1.
	Extracti Hyoscyami Viride	.. gr. 2.

Mix. Varnish. Dose, 1.

General Formulæ.

PILULA ALOES CUM BELLADONNÂ.

℞ Extracti Aloës Barbadensis .. gr. 1.
Jalapæ Resinæ .. .. .. gr. $\frac{1}{2}$.
Extracti Belladonnæ Viride .. gr. $\frac{1}{4}$.
Confectionem Rosæ Gallicæ ad gr. 3.
Mix. Varnish. Dose, 1.

PILULA ATROPINÆ SULPHATIS.

℞ Atropinæ Sulphatis .. .. gr. $\frac{1}{120}$.
Pulveris Glycyrrhizæ .. .. gr. 2.
Pulveris Tragacanthæ .. .. gr. 1.
Aquæ .. .. q. s.
Mix. Dose, 1 or 2.

PILULA CALOMELANOS CUM COLOCYNTHIDE.

℞ Hydrargyri Subchloridi .. gr. 1.
Pulveris Ipecacuanhæ .. .. gr. $\frac{1}{2}$.
Extracti Colocynthidis Compositi .. .. gr. $3\frac{1}{2}$.
Mix. Dose, 1 or 2.

PILULA COLCHICI COMPOSITA.

℞ Extracti Colchici.. .. .. gr. 1.
Pulveris Ipecacuanhæ Compositi gr. 3.
Mix. Dose, 1.

General Formulæ.

PILULA COLOCYNTHIDIS CUM HYDRARGYRO.

℞ Pilulæ Hydrargyri
Extracti Colocynthidis Compositi, āā .. .. .. gr. $2\frac{1}{2}$.

Mix. Dose, 1 or 2.

PILULA CUPRI COMPOSITA.

℞ Cupri Sulphatis .. .. .. gr. $\frac{1}{4}$.
Pulveris Opii .. .. .. gr. $\frac{1}{2}$.
Extractum Gentianæ .. ad gr. 3.

Mix. Varnish with Keratin. Dose, 1.

PILULA DIGITALIS COMPOSITA.

℞ Pulveris Digitalis .. .. gr. $\frac{1}{2}$.
Pulveris Scillæ
Pilulæ Hydrargyri, āā .. .. gr. 1.
Theriacæ .. .. q. s.

Mix. Dose, 1.

PILULA DIGITALIS CUM SCILLÂ.

℞ Pulveris Digitalis .. .. gr. $\frac{1}{2}$.
Pilulæ Scillæ Compositæ .. gr. $4\frac{1}{2}$.

Mix. Dose, 1.

Sir Wm Gowers for Trigeminal Neuralgia

Liq. Trinitrini m p.
Inf. Gelsemii ℥ xii
Liq. Strychninae m V
Syr. Aurantii ℥ ℈
Aquam ad ℥ ii

R.d.s

General Formulæ.

PILULA ELATERINI COMPOSITA.

℞ Elaterini .. .. gr. $\frac{1}{20}$.
Extracti Aloës Barbadensis .. gr. 1.
Extracti Hyoscyami Viride .. gr. 2.
Mix. Dose, 1.

PILULA FERRI SULPHATIS EXSICCATÆ.

℞ Ferri Sulphatis Exsiccatæ .. gr. 2.
Theriacæ .. .. q. s.
Mix. Varnish. Dose, 1.

PILULA FERRI SULPHATIS COMPOSITA.

℞ Ferri Sulphatis Exsiccatæ
Quininæ Sulphatis, āā .. .. gr. 1.
Extracti Aloës Barbadensis .. gr. $\frac{1}{2}$.
Extracti Belladonnæ Viride
Extracti Nucis Vomicæ, āā .. gr. $\frac{1}{4}$.
Mix. Varnish. Dose, 1 or 2.

PILULA HYDRARGYRI ET ALOËS.

℞ Pilulæ Hydrargyri
Extracti Aloës Barbadensis, āā.. gr. $1\frac{1}{2}$.
Extracti Hyoscyami Viride .. gr. 2.
Mix. Dose, 1 or 2.

General Formulæ.

PILULA HYDRARGYRI CUM RHEO.

℞ Pilulæ Hydrargyri .. .. gr. 1½.
Pulveris Rhei .. .. .. gr. 3.
Pulveris Ipecacuanhæ .. .. gr. ½.
Theriacæ .. .. q. s.
Mix. Dose, 1 or 2.

PILULA QUININÆ CUM FERRO.

℞ Quininæ Sulphatis
Ferri Sulphatis, āā .. .. gr. 1.
Confectionis Rosæ Gallicæ .. q. s.
Mix. Varnish. Dose, 1.

PILULA QUININÆ CUM OPIO.

℞ Quininæ Sulphatis .. .. gr. 1.
Pulveris Opii .. .. .. gr. ¼.
Confectionis Rosæ Gallicæ .. q. s.
Mix. Dose, 1.

PILULA ZINCI OXIDI CUM BELLADONNÂ.

℞ Zinci Oxidi .. .. .. gr. 2.
Extracti Belladonnæ Viride .. gr. ¼.
Mix. Dose, 1.

General Formulæ.

PULVIS MAGNESII SULPHATIS COMPOSITUS.

℞ Magnesii Sulphatis .. .. .. gr. 60.
Sodii Sulphatis .. .. .. gr. 30.
Magnesii Carbonatis Levis .. gr. 5.
Pulveris Zingiberis .. .. gr. 3.

Mix.

PULVIS STRAMONII COMPOSITUS.

℞ Pulveris Stramonii .. .. oz. 1.
Pulveris Anisi Fructûs
Potassii Nitratis, āā .. .. oz. ½.
Tabaci Contriti .. .. .. gr. 30.

Mix. Burn as much as will lie on a shilling and inhale the smoke.

SUPPOSITORIUM ACIDI TANNICI CUM MORPHINÂ.

℞ Acidi Tannici .. .. .. gr. 5.
Morphinæ Hydrochloridi .. gr. ¼.
Oleum Theobromatis .. ad gr. 20.

Mix.

SUPPOSITORIUM BELLADONNÆ.

℞ Extracti Belladonnæ Viride .. gr. ½.
Oleum Theobromatis .. ad gr. 20.

General Formulæ.

SUPPOSITORIUM BELLADONNÆ CUM MORPHINÂ.

℞ Extracti Belladonnæ Viride .. gr. ½.
Morphinæ Hydrochloridi .. gr. ¼.
Oleum Theobromatis .. ad gr. 20.
Mix.

SUPPOSITORIUM GALLÆ COMPOSITUM.

℞ Pulveris Gallæ .. .. .. gr. 3.
Pulveris Opii .. .. .. gr. 1.
Oleum Theobromatis .. ad gr. 20.
Mix.

SUPPOSITORIUM NUTRIENS (A).

℞ Peptonized Milk Concentrated
Olei Theobromatis, āā .. .. gr. 20.
Mix.

SUPPOSITORIUM NUTRIENS (B).

℞ Peptonized Beef (Pepsin Peptone)
Olei Theobromatis, āā .. .. gr. 20.
Mix.

SUPPOSITORIUM NUTRIENS (C).

℞ Peptonized Cocoa and Milk Concentrated
Olei Theobromatis, āā .. .. gr. 20.
Mix.

General Formulæ.

SUPPOSITORIUM OPII.

℞ Pulveris Opii .. .. .. gr. 1.
Oleum Theobromatis .. ad gr. 20.

Mix.

UNGUENTUM BELLADONNÆ COMPOSITUM.

℞ Extracti Belladonnæ Viride
Unguenti Potassii Iodidi, āā .. oz. 1.

Mix.

UNGUENTUM HYDRARGYRI CUM BELLADONNÂ.

℞ Unguenti Belladonnæ Viride
Unguenti Hydrargyri, āā .. oz. 1.

Mix.

UNGUENTUM HYDRARGYRI CUM OPIO.

℞ Pulveris Opii .. .. .. oz. $\frac{1}{4}$.
Unguenti Hydrargyri .. .. oz. 2.

Mix.

General Formulæ.

UNGUENTUM HYDRARGYRI MITIUS.

℞ Unguenti Hydrargyri
Adipis Lanæ
Paraffini mollis, āā .. .. oz. 1.

Mix.

UNGUENTUM HYDRARGYRI NITRATIS MITIUS.

℞ Unguenti Hydrargyri Nitratis oz. ¼.
Paraffini mollis .. .. .. oz. 2.

Mix.

UNGUENTUM HYDRARGYRI OXIDI RUBRI MITIUS.

℞ Unguenti Hydrargyri Oxidi
Rubri .. .. oz. ¼.
Adipis Benzoati .. .. .. oz. 1.

Mix.

UNGUENTUM OPII.

℞ Pulveris Opii .. .. .. oz. ¼.
Adipis Benzoati .. .. .. oz. 2.

Mix.

General Formulæ.

UNGUENTUM SIMPLEX.

℞ Ceræ Flavæ
Adipis Benzoati, āā .. .. oz. 1.
Olei Olivæ .. .. .. fl. oz. iv.

Mix.

UNGUENTUM STRAMONII.

℞ Foliorum Stramonii Recentium lb. ½.
Adipis Benzoati .. .. .. lbs. 2.

Mix the bruised leaves with the lard, and expose to a mild heat until the leaves become friable; strain through linen.

B.

FORMULÆ FOR CHILDREN.

*** *The doses here given are intended for children up to two years of age.*

MISTURA ACIDI HYDROCHLORICI.

℞ Acidi Hydrochlorici diluti
Spiritûs Chloroformi
Syrupi, āā .. .. .. fl. dr. ss.
Aquam destillatam .. ad fl. oz. iss.

Mix. Dose, fl. dr. j. to fl. dr. ij.

MISTURA ALUMINIS.

℞ Aluminis .. .. gr. 40.
Syrupi Rhœados.. .. .. fl. dr. j.
Aquam destillatam .. ad. fl. oz. iss.

Mix. Dose, fl. dr. j.

MISTURA AMMONII ACETATIS.

℞ Liquoris Ammonii Acetatis .. fl. dr. iij.
Potassii Citratis .. .. .. gr. 60.
Glycerini .. .. fl. dr. j.
Aquam destillatam .. ad fl. oz. iss.

Mix. Dose, fl. dr. j. to fl. dr. ij.

Asthma in Children

R. a. Y.

Inst Hosp. Bür

Sol. Iod 8 ii

D. Lobelinae Alk gtt iij

Formulæ for Children.

MISTURA ÆTHERIS CUM AMMONIÂ.

℞ Ammonii Carbonatis .. .. gr. 3.
Syrupi Scillæ .. .. .. fl. dr. iss.
Haustum Ætheris Compositum .. .. .. ad fl. oz. iss.

Mix. Dose, fl. dr. j. to fl. dr. ij.

MISTURA AROMATICA.

℞ Magnesii Carbonatis Levis .. gr. 12.
Pulveris Cretæ Aromatici .. gr. 60.
Tincturæ Rhei Compositæ .. fl. dr. j.
Aquam Menthæ Piperitæ ad fl. oz. iss.

Mix. Dose, fl. dr. j. to fl. dr. ij.

MISTURA CARMINATIVA.

℞ Sodii Bicarbonatis .. .. gr. 24.
Spiritûs Ammoniæ Aromatici .. min. xxiv.
Glycerini .. .. fl. dr. j.
Syrupi .. .. fl. dr. ij.
Aquam Menthæ Piperitæ ad fl. oz. iss.

Mix. Dose, fl. dr. j. to fl. dr. ij.

Formulæ for Children.

MISTURA CHLORAL.

℞ Chloral Hydratis
Potassii Bromidi, āā .. .. gr. 30.
Syrupi Tolutani .. .. .. fl. dr. iss.
Aquam Cassiæ .. .. ad fl. oz. iss.

Mix. Dose, fl. dr. j. to fl. dr. ij.

MISTURA CINCHONÆ.

℞ Acidi Nitrici diluti .. .. min. xxx.
Tincturæ Cinchonæ .. .. fl. dr. ij.
Glycerini .. .. fl. dr. j.
Aquam destillatam .. ad fl. oz. iss.

Mix. Dose, fl. dr. j. to fl. dr. ij.

MISTURA HÆMATOXYLI.

℞ *Extracti Hæmatoxyli* .. .. gr. 60.
Tincturæ Catechu .. .. fl. dr. ij.
Aquæ ferventis
Aquæ Carui, āā .. .. .. fl. dr. v.

Mix. Dose, fl. dr. j. to fl. dr. ij.

MISTURA HYDRARGYRI PERCHLORIDI.

℞ Liquoris Hydrargyri Perchloridi min. xxiv.
Glycerini .. .. fl. dr. ij.
Aquam Carui .. .. ad fl. oz. iss.

Mix. Dose, fl. dr. ss. to fl. dr. j.

Formulæ for Children.

MISTURA IPECACUANHÆ.

℞	Vini Ipecacuanhæ ..	.. fl. dr. ss.
	Tincturæ Scillæ	.. min. xxx.
	Glycerini	.. fl. dr. j.
	Aquam destillatam ..	ad fl. oz. iss.

Mix. Dose, fl. dr. j. to fl. dr. ij.

MISTURA IPECACUANHÆ AMMONIATA.

℞	Vini Ipecacuanhæ ..	.. fl. dr. ss.
	Tincturæ Scillæ	.. min. xxx.
	Spiritûs Ammoniæ Aromatici	.. min. xxiv.
	Glycerini	.. fl. dr. j.
	Aquam destillatam ..	ad fl. oz. iss.

Mix. Dose, fl. dr. j. to fl. dr. ij.

MISTURA IPECACUANHÆ OPIATA.

℞	Sodii Bicarbonatis ..	.. gr. 20.
	Spiritûs Ætheris Nitrosi	.. min. xxx.
	Tincturæ Opii	.. min. iij.
	Misturam Ipecacuanhæ ..	ad fl. oz. iss.

Mix. Dose, fl. dr. j. to fl. dr. ij.

Formulæ for Children.

MISTURA OLEI MORRHUÆ CUM FERRO.

℞ Olei Morrhuæ
Vini Ferri, āā .. .. .. fl. oz. ss.
Olei Cassiæ .. .. .. min. ij.
Mucilaginis Acaciæ .. .. fl. dr. ij.

Mix. Dose, fl. dr. j. to fl. dr. ij.

MISTURA OLEI RICINI.

℞ Olei Ricini .. .. .. fl. dr. iss.
Pulveris Acaciæ .. .. .. gr. 30.
Syrupi Simplicis .. .. .. fl. dr. j.
Aquam Carui .. .. ad fl. oz. iss.

Mix. Dose, fl. dr. j. to fl. dr. ij.

PULVIS HYDRARGYRI CUM MAGNESIÂ.

℞ Hydrargyri cum Cretâ
Pulveris Rhei
Pulveris Cinnamomi, āā .. gr. 1.
Magnesii Carbonatis Levis .. gr. 2.

Mix.

PULVIS RHEI CUM CALOMELANE.

℞ Pulveris Rhei .. .. .. gr. 4.
Hydrargyri Subchloridi
Pulveris Cretæ Aromatici, āā .. gr. 1.

Mix.

Formulæ for Children.

PULVIS RHEI CUM HYDRARGYRO.

℞ Pulveris Rhei .. .. .. gr. 3.
Hydrargyri cum Cretâ .. .. gr. 2.
Mix.

PULVIS RHEI CUM JALAPÂ.

℞ Pulveris Rhei .. .. .. gr. 1.
Pulveris Jalapæ Compositi .. gr. 2.
Mix.

PULVIS RHEI CUM SODÂ.

℞ Pulveris Rhei .. .. .. gr. 6.
Sodii Bicarbonatis .. .. gr. 3.
Pulveris Zingiberis .. .. gr. 1.
Mix.

PULVIS SCAMMONII ET CALOMELANOS CUM JALAPÂ.

℞ Hydrargyri Subchloridi.. .. gr. 1.
Pulveris Scammonii Resinæ .. gr. 1.
Pulveris Jalapæ .. .. .. gr. 4.
Mix.

PULVIS SCAMMONII CUM CALOMELANE.

℞ Pulveris Scammonii Resinæ .. gr. 3.
Hydrargyri Subchloridi
Sacchari Purificati, āā .. .. gr. 2.
Mix.

C.

FORMULÆ FOR THE EYE DEPARTMENT.

COLLYRIUM ACIDI BORICI ET ZINCI SULPHATIS.

℞	Acidi Borici	.. gr. 5.
	Zinci Sulphatis	.. gr. ½.
	Aquæ destillatæ	.. fl. oz. j.

Dissolve.

COLLYRIUM ARGENTI NITRATIS.

℞	Argenti Nitratis.. ..	.. gr. 1.
	Aquæ destillatæ	.. fl. oz. j.

Dissolve.

COLLYRIUM BORACIS.

℞	Boracis	.. gr. 10.
	Aquæ destillatæ	.. fl. oz. j.

Dissolve.

Formulæ for the Eye Department.

COLLYRIUM ZINCI SULPHATIS.

℞ Zinci Sulphatis .. .. .. gr. 1.
Aquæ destillatæ .. .. .. fl. oz. j.

Dissolve.

GUTTÆ ATROPINÆ.

℞ Atropinæ Sulphatis .. .. gr. 1.
Aquæ destillatæ .. .. .. fl. oz. j.

Dissolve.
If a stronger preparation is required, the strength must be stated.

GUTTÆ CUPRI SULPHATIS.

℞ Cupri Sulphatis .. .. .. gr. 1.
Aquæ destillatæ .. .. .. fl. oz. j.

Dissolve.

GUTTÆ PHYSOSTIGMINÆ.

℞ Physostigminæ Sulphatis .. gr. 1.
Aquæ destillatæ .. .. .. fl. oz. j.

Dissolve.

Formulæ for the Eye Department.

GUTTÆ ZINCI CHLORIDI.

℞ Zinci Chloridi .. .. .. gr. 1.
Aquæ destillatæ .. .. .. fl. oz. j.

Dissolve.

UNGUENTUM ATROPINÆ.

℞ Atropinæ.. .. .. .. gr. 4.
Vaselini .. .. oz. j.

Mix.

UNGUENTUM HYDRARGYRI OXIDI FLAVI.

℞ Hydrargyri Oxidi Flavi .. gr. 8.
Vaselini .. .. oz. j.

Mix.

UNGUENTUM HYDRARGYRI OXIDI FLAVI CUM ATROPINÂ.

℞ Hydrargyri Oxidi Flavi .. gr. 8.
Atropinæ.. .. .. .. gr. 4.
Vaselini .. .. oz. j.

Mix.

UNGUENTUM HYDRARGYRI OXIDI FLAVI CUM PHYSOSTIGMINÂ.

℞ Hydrargyri Oxidi Flavi .. gr. 8.
Physostigminæ .. .. .. gr. ½.
Vaselini .. .. oz. j.

D.

FORMULÆ FOR THE EAR AND THROAT DEPARTMENT.

GARGARISMA ACIDI TANNICI.

℞ Glycerini Acidi Tannici .. fl. oz. ss.
Aquam destillatam .. ad fl. oz. x.

Mix.

GARGARISMA ALUMINIS.

℞ Aluminis .. .. gr. 120.
Glycerini .. .. fl. oz. ss.
Aquam destillatam .. ad fl. oz. xij.

Mix.

GARGARISMA BORACIS.

℞ Boracis .. .. oz. ½.
Oxymellis .. .. .. fl. oz. j.
Aquam destillatam .. ad fl. oz. xij.

Mix.

Formulæ for the Ear and Throat Department.

GARGARISMA CHLORI.

℞ *Liquoris Chlori* ..	..	.. fl. oz. j.
Aquam destillatam	..	ad fl. oz. xij.

Mix.

GARGARISMA CUPRI SULPHATIS.

℞ Cupri Sulphatis ..	..	.. gr. 48.
Glycerini	..	.. fl. oz. j.
Aquam destillatam	..	ad fl. oz. xij.

Mix.

GARGARISMA HYDRARGYRI PERCHLORIDI.

℞ Hydrargyri Perchloridi..		.. gr. 2.
Acidi Hydrochlorici diluti		.. fl. dr. ij.
Glycerini	..	.. fl. oz. j.
Aquam destillatam	..	ad fl. oz. x.

Mix.

GARGARISMA MYRRHÆ.

℞ Tincturæ Myrrhæ	..	.. fl. oz. ss.
Aquam destillatam	..	ad fl. oz. xij.

Mix.

Formulæ for the Ear and Throat Department.

GARGARISMA POTASSII CHLORATIS.

℞ Potassii Chloratis .. .. oz. ½.
Aquæ destillatæ .. .. .. fl. oz. xij.

Dissolve.

GARGARISMA SODII CHLORINATÆ.

℞ Liquoris Sodæ Chlorinatæ .. fl. oz. ss.
Aquam destillatam .. ad fl. oz. xij.

Mix.

INJECTIO ACIDI CARBOLICI COMPOSITA.

℞ Acidi Carbolici .. .. .. gr. 80.
Acidi Borici .. .. .. gr. 120.
Aquæ destillatæ .. .. .. fl. oz. x.

Dissolve.
An injection for the ear.

INJECTIO BORACIS COMPOSITA.

℞ Boracis .. .. gr. 240.
Sodii Chloridi .. .. .. gr. 60.
Aquæ destillatæ .. .. fl. oz. x.

Dissolve.
A nasal douche. Add fl. oz. ss. to water O j.

Formulæ for the Ear and Throat Department.

PIGMENTUM FERRI.

℞ Liquoris Ferri Perchloridi .. fl. dr. j.
Glycerinum .. .. ad fl. oz. j.

Mix.

PULVIS BISMUTHI COMPOSITUS.

℞ Bismuthi Subnitratis .. .. gr. 20.
Pulveris Acidi Tannici .. .. gr. 10.
Quininæ Sulphatis .. .. gr. 3.
Pulveris Amyli .. .. .. oz. 1.

Mix. To be used as a snuff.

Inst Capsici Subsulphas

℞ Tinct Capsici ℳ v
Tinct Zingiberis ℳ xx
Potass Bromidi gr x
Sodii Bicarbonatis gr x
Decoct Cinchonae ℨj/s.

E.

FORMULÆ FOR THE SKIN DEPARTMENT.

BALNEA (BATHS).

The quantity given is that suitable for a full bath of 30 gallons of water, usually at a temperature of 90—95° Fahr.

BALNEUM ALKALINUM.

℞ Potassii Carbonatis
Sodii Carbonatis, āā .. .. oz. 3.

Used for itchy eczema, urticarial and pruriginous conditions, and for removing the scales of psoriasis. The bath should be prolonged for 20 minutes to half an hour.

BALNEUM IODI.

℞ Iodi .. .. gr. 30.
Potassii Iodidi .. .. .. oz. ½.
Glycerini .. .. fl. oz. ij.

Used in tuberculous, scrofulous, and late syphilitic affections of the skin.

Formulæ for the Skin Department.

BALNEUM MERCURIALE.

℞ Hydrargyri Perchloridi.. .. gr. 120.
Acidi Hydrochlorici diluti .. fl. dr. j.

Used for early rupioid syphilides.

BALNEUM SULPHURIS COMPOSITUM.

℞ Sulphuris Precipitati .. .. oz. 2.
Sodii Hyposulphitis .. .. oz. 1.
Acidi Sulphurici diluti .. .. fl. oz. ss.
Aquam .. .. .. ad O j.

Used in the treatment of scabies in children or in persons specially liable to irritation from sulphur ointment.

GLYCOGELATINA ZINCI OXIDI.

℞ Zinci Oxidi
Gelatini, āā .. .. .. gr. 90.
Glycerini.. .. .. .. fl. dr. iij.
Aquæ destillatæ .. .. .. fl. dr. iv.

Directions.—Rub the oxide into a smooth paste with the glycerine; then mix with the gelatine previously dissolved

in hot water and stir till cold. To be melted over a water bath or in a glue pot, and applied with a flat brush, then dabbed with absorbent wool to prevent sticking to the clothes.

An application employed for many forms of chronic dermatitis with little or no discharge, especially upon the lower extremities. It is also used as an anti-pruritic in lichenous conditions, but is not generally applicable to hairy parts, as its removal entails pain. The preparation is usually stripped off at the end of a week.

LINIMENTUM CALAMINÆ.

℞	Calaminæ Præparatæ ..	.. gr. 40.
	Zinci Oxidi	.. gr. 30.
	Aquæ Calcis	
	Olei Olivæ, āā	.. fl. oz. ss.

The affected part to be wrapped in soft linen soaked in the liniment.

Frequent moistening or renewal of the liniment is necessary.

A preparation applicable for all forms of acute dermatitis.

Formulæ for the Skin Department.

LOTIO ACIDI SALICYLICI CUM BORACE.

℞	Acidi Salicylici	.. gr. 120.
	Boracis	.. gr. 60.
	Glycerini	
	Spiritûs Rectificati, āā ..	.. fl. oz. j.
	Aquam destillatam ..	ad fl. oz. x.

Used in urticaria, prurigo, and itching affections.

LOTIO ACIDI SULPHUROSI.

℞	Acidi Sulphurosi ..	.. fl. oz. j.
	Aquam destillatam ..	ad fl. oz. viij.

Used in tinea versicolor.

LOTIO AMMONIÆ ET CANTHARIDIS.

℞	Liquoris Ammoniæ Fortioris	.. fl. dr. j.
	Spiritûs Rosmarini ..	.. fl. dr. ij.
	Spiritûs Rectificati ..	.. fl. dr. iv.
	Tincturæ Cantharidis ..	.. fl. dr. ij.
	Olei Amygdalæ	
	Olei Olivæ, āā	.. fl. oz. j.

Used in alopecia when no seborrhœa or inflammatory condition is present.

Formulæ for the Skin Department.

LOTIO BISMUTHI.

℞ Bismuthi Subnitratis .. .. gr. 8.
Zinci Oxidi .. .. .. gr. 30.
Hydrargyri Perchloridi .. gr. ¼.
Glycerini .. .. min. xv.
Aquam destillatam .. ad fl. oz. j.

A stimulant and cosmetic application used in rosacea.

LOTIO CALAMINÆ.

℞ Calaminæ .. .. .. gr. 30.
Zinci Oxidi .. .. .. gr. 20.
Glycerini .. .. min. xv.
Aquæ destillatæ .. .. .. fl. oz. j.

Used in rosacea, lupus, erythematosus, and erythema.

LOTIO CARBONIS DETERGENS.

℞ Liquoris Carbonis Detergentis
fl. dr. j. to fl. dr. ij.
Spiritûs Rectificati .. .. fl. dr. iij.
Aquam destillatam .. ad fl. oz. viij.

An astringent, and anti-pruritic evaporating lotion, used in hyperæmic and itching conditions; must be used with caution in eczemas.

Formulæ for the Skin Department.

LOTIO PARAFFINI COMPOSITA.

℞ Paraffini .. .. oz. 3.
Balsami Peruviani .. .. fl. dr. ij.
Hydrargyri Oleatis .. .. gr. 60.
Olei Olivæ .. .. .. fl. oz. iss.

To be applied to the head with a stiff brush ("sash-tool") every two hours for twelve hours. Destroys pediculi, and dislodges their nits.

LOTIO PLUMBI CUM PICE.

℞ Liquoris Plumbi Subacetatis
fl. dr. j. to fl. dr. ij.
Liquoris Picis Carbonis fl. dr. j. to fl. dr. ij.
Aquam destillatam .. ad fl. oz. viij.

An anti-pruritic in urticaria, etc.

LOTIO PLUMBI LACTATIS.

℞ Liquoris Plumbi Subacetatis .. fl. dr. j.
Acidi Salicylici .. .. .. gr. 2.
Lac .. .. .. ad fl. oz. ij.

Used in erythema and many phases of eczema, with or without discharge.

Formulæ for the Skin Department.

LOTIO SAPONIS MOLLIS.

℞	Saponis Mollis	.. oz. 1.
	Olei Lavandulæ	.. min. xx.
	Spiritûs Rectificati ..	.. fl. oz. iss.
	Aquam destillatam ..	ad fl. oz. viij.

A stimulant in dry chronic eczematoid conditions, or as a scalp wash in seborrhœa and psoriasis.

LOTIO SODII HYPOSULPHITIS.

℞	Sodii Hyposulphitis ..	.. gr. 300.
	Acidi Sulphurosi.. ..	.. fl. oz. j. ss.
	Glycerini..	.. fl. oz. j.
	Aquam destillatam ..	ad O j.

Used in tinea versicolor.

LOTIO SULPHURIS.

℞	Sulphuris Præcipitati ..	.. gr. 240.
	Spiritûs Rectificati ..	.. fl. oz. ss.
	Glycerini..	.. fl. dr. ij.
	Aquæ destillatæ	.. fl. oz. iv.

Used in acne, vulgaris, and rosacea, and in seborrhœic disorders.

Formulæ for the Skin Department.

LOTIO SULPHURIS COMPOSITA.

℞	Sulphuris Præcipitati	
	Potassii Carbonatis, āā ..	.. gr. 120.
	Spiritûs Rectificati	
	Ætheris	
	Glycerini, **āā**	.. fl. dr. ij.
	Aquam destillatam ..	ad fl. oz. viij.

Employed in seborrhœic disorders with dermatitis, scabies, and tinea versicolor.

PASTA BARII SULPHIDI.

℞	Barii Sulphidi	.. gr. 120.
	Zinci Oxidi	
	Pulveris Amyli, **āā** ..	.. gr. 180.

To be made into a paste with water, and applied for 10 to 15 minutes.

A depilatory, used as a preliminary treatment in "sycosis." Must be watched with care, as it is apt to cause severe dermatitis.

Ichthyol after food
does away with the flushing after food.

Formulæ for the Skin Department.

PASTA ZINCI CUM SULPHURE.

℞ Unguenti Zinci
Unguenti Sulphuris, āā .. oz. ½.
Pulveris Amyli .. .. gr. 60 to gr. 120.

A gentle stimulant in acne and rosacea. Its consistence renders it pleasanter for application than ordinary ointments. Is best applied at night after vigorous washing.

PASTA IODI ET PICIS. (Coster's Paste.)

℞ Iodi .. .. gr. 120.
Olei Picis Liquidi .. .. fl. oz. j.

To be applied with a stiff brush.

Used in obstinate, chronic, localized ringworm. Must be applied with caution, as kerion is apt to supervene upon its employment.

PASTA ZINCI OXIDI CUM ACIDO SALICYLICO. (Lassar's Paste.)

℞ Zinci Oxidi
Pulveris Amyli, āā .. .. gr. 120.
Vaselini .. .. oz. ½.
Acidi Salicylici .. .. .. gr. 10.

To be applied thickly and covered with butter-cloth.

Used in various eczematoid conditions with moderate amount of discharge.

Formulæ for the Skin Department.

PASTA ZINCI OXIDI CUM RESORCIN.
(Ihle's Paste.)

℞ Zinci Oxidi
Vaselini
Adipis Lanæ
Pulveris Amyli, āā .. .. gr. 120.
Resorcini.. .. .. .. gr. 10.

Very similar in properties to Lassar's paste, but rather more stimulating.

PIGMENTUM ACIDI SALICYLICI.

℞ Acidi Salicylici .. .. .. gr. 60.
Extracti Cannabis Indicæ .. gr. 10.
Cocainæ Hydrochloridi .. gr. 5.
Collodium flexile .. ad fl. oz. j.

Used as a solvent for corns, callosities, and other forms of hyperkeratosis.

PIGMENTUM CHRYSAROBINI.

℞ Gutta Perchæ .. .. .. gr. 60.
Chloroformi .. .. .. dr. x.
Chrysarobini .. .. .. gr. 60.

To be painted on with a strong brush.

Used in psoriasis as limiting the action of the chrysarobin to spots affected, and in diminishing the amount of staining of clothes, etc.

Formulæ for the Skin Department.

SPIRITÛS SAPONIS ALKALINUS.
(Hebra.)

℞ Saponis Mollis .. .. .. oz. 4.
Spiritûs Rectificati .. .. fl. oz. ij.

A shampooing liquid, used in many scaly conditions of the scalp.

UNGUENTUM BALSAMI PERUVIANI.

℞ Balsami Peruviani .. .. fl. dr. j.
Adipis Præparati .. .. oz. 1.

Used in scabies for persons who cannot use sulphur.

UNGUENTUM CHRYSAROBINI COMPOSITUM.

℞ Chrysarobini .. .. .. gr. 150.
Ichthyol .. .. gr. 150.
Acidi Salicylici .. .. .. gr. 60.
Paraffinum Molle .. ad oz. 8.

Used in chronic diffuse ringworm of the scalp, and occasionally in persistent seborrhœic eczema of the flexures. In certain individuals it causes erysipelatoid dermatitis.

Formulæ for the Skin Department.

UNGUENTUM CRETÆ.

℞	Cretæ Præparatæ ..	.. oz. $1\frac{1}{2}$.
	Adipis Benzoati	.. oz. 1.

Used in erythema and eczematoris conditions, especially of the face.

UNGUENTUM DIACHYLI.

℞	Emplastri Plumbi ..	.. oz. 2.
	Olei Olivæ	.. fl. oz. iij.

Used in chronic eczema with much infiltration.

UNGUENTUM HYDRARGYRI CUM ACIDO CARBOLICO.

℞	Hydrargyri Perchloridi ..	.. gr. 3.
	Acidi Carbolici	.. gr. 20.
	Olei Olivæ	.. min. xx.
	Unguentum Zinci ..	.. oz. 1.

Used in itching dry dermatitis, lichen planus, and impetigo.

Formulæ for the Skin Department.

UNGUENTUM HYDRARGYRI NITRATIS COMPOSITUM.

℞ Acidi Carbolici
Unguenti Hydrargyri Nitratis
Unguenti Sulphuris, āā .. .. oz. 1.

Used in chronic ringworm; also in impetigo and pustular sycosis.

UNGUENTUM METALLORUM.

℞ Unguenti Hydrargyri Nitratis diluti
Unguenti Plumbi Subacetatis
Unguenti Zinci, āā .. .. oz. 1.

Used for dry eczema, especially of the palms and soles.

UNGUENTUM NAPHTHOL COMPOSITUM.

℞ Naphthol	.. ..	gr. 15 to gr. 60.
Cretæ Preparatæ	.. ..	gr. 20 to gr. 80.
Saponis Mollis	.. ..	gr. 120 to oz. 1.
Adipis Benzoatis	.. ..	gr. 180 to oz. $1\frac{1}{2}$.

Used for scabies where sulphur applications are contra-indicated.

Formulæ for the Skin Department.

UNGUENTUM PYROGALLOL COMPOSITUM.

℞ Acidi Pyrogallici .. .. gr. 30 to gr. 60.
Ichthyol .. .. .. gr. 30.
Acidi Salicylici .. .. gr. 15.
Paraffinum Molle .. ad oz. 1.

Used in psoriasis where chrysarobin preparations fail, and in lupus vulgaris.

UNGUENTUM ZINCI CUM ACIDO SALICYLICO.

℞ Acidi Salicylici .. .. .. gr. 20.
Unguenti Zinci
Paraffini Mollis, āā .. .. oz. ½.

Used in eczematoid conditions with small amount of discharge.

TREATMENT OF POISONING.

Remarks.

If the symptoms point to the conclusion that some powerful Irritant or Corrosive has been swallowed, it will be well to administer at once, without waiting to obtain more precise information, a mixture of Carron Oil or Milk, with Chalk and White of Egg, such as the following :—

℞	Linimenti Calcis	..	.. fl. oz. iij.
	Cretæ Præparatæ	..	.. gr. 120.
	Albuminis Ovi ..	..	.. fl. oz. vj.

Mix.

Albumen forms insoluble compounds with nearly all the metallic salts, generally combining with both acid and base ; it will therefore be improper to administer emetics such as Sulphate of Zinc or Sulphate of Copper, when Albumen has been previously given as an antidote ; in this

case vomiting should be induced by means of Mustard and Warm Water, or Ipecacuanha Wine.

Vinegar increases the solubility of most poisons; it should therefore never be given as an antidote, except, perhaps, in a case of poisoning by the Caustic Alkalies, and even then Oil is better.

It is of the greatest importance to treat the condition of collapse which so often accompanies cases of poisoning. This is best accomplished by placing the patient in a warm bed, with hot-water bottles to the extremities, and administering an enema of an ounce of brandy with two ounces of warm water or beef-tea.

ANTIDOTES, ETC.

ACID CARBOLIC.

Olive and Castor Oil, Stomach Pump, or Emetics.

Strong Tea or Coffee to counteract narcotism.

ACID HYDROCYANIC, OR CYANIDE OF POTASSIUM.

A mixture of Proto- and Per-Salt of Iron, with an Alkaline Carbonate.

The Inhalation of *Diluted* Ammonia or Chlorine.

Cold Affusion to head and spine.

℞	Ferri Sulphatis	.. gr. 10.
	Liquoris Ferri Perchloridi	.. fl. dr. j.
	Aquam	ad fl. oz. j.

Mix. Administer this draught, and immediately afterwards give 20 grains of Carbonate of Potassium dissolved in an ounce of Water.

This will render insoluble 2 grains of pure Hydrocyanic Acid (=110 minims of the dilute Acid of the Pharmacopœia) or 5 grains of Cyanide of Potassium.

Stomach Pump, Artificial Respiration, and Galvanism.

ACIDS MINERAL.

Calcined Magnesia or Chalk, with White of Egg, Carron Oil, or Soap-suds, followed by Enemata of Beef Tea and Brandy with Tincture of Opium to prevent collapse.

ACID OXALIC, AND SOLUBLE OXALATES.

Chalk, Emetics, Stomach Pump, Enemata of Beef Tea with Brandy and Opium to prevent collapse.

NOTE.—The use of the Stomach Pump is attended with some danger in cases where there is much erosion, especially if there be hæmorrhage from the stomach.

ALCOHOL, ETHER, CHLOROFORM, CHLORAL HYDRATE, AND THE HYDROCARBONS.

Stomach Pump, Emetics, Cold Affusion to head and spine.

Electro-Magnetism, Artificial Respiration.

CAUSTIC ALKALIES.

Olive Oil, Dilute Acetic Acid or Vinegar.

NOTE.—This treatment is also suitable for poisoning by Ammoniated Camphor Liniment.

ALKALOIDS.

(ACONITINE, ATROPINE, STRYCHNINE.)

Emetics and Stomach Pump. No reliable antidote.

Chloral Hydrate has proved useful in some cases of poisoning by Strychnine.

TARTARATED ANTIMONY.

(Tartar Emetic.)

Tannic Acid, Catechu, Decoction of Yellow Bark or other Vegetable Astringents. Recently precipitated Peroxide of Iron (see Arsenic).

ANTIMONY, CHLORIDE OF.

(Butter of Antimony.)

Magnesia, Carbonate of Sodium.

ARSENIC AND ARSENICAL COMPOUNDS.

Recently precipitated Peroxide of Iron, Carron Oil, Emetics, Stomach Pump, Artificial Respiration.

℞ Liquoris Ferri Perchloridi .. fl. oz. x.
Sodii Carbonatis.. .. .. oz. 4.
Aquam .. .. .. ad O ij.

Mix.

One fluid ounce of this mixture will render 1 grain of Arsenic insoluble.

BARIUM, SOLUBLE SALTS OF.

Sulphate of Magnesium or Sulphate of Sodium.

BELLADONNA AND ITS SALTS.

Stomach Pump and Emetics.

AMMONIATED CAMPHOR LINIMENT.

See Note to Caustic Alkalies.

CANTHARIDES.

Emetics or Stomach Pump, followed by Castor Oil and Opium. There is no antidote, but Carron Oil and Albumen may be given when the Poison has been removed.

COPPER, SOLUBLE SALTS OF.

Emetics, Albumen, Iron Filings.

IODINE.

Albumen or Milk, with Flour and Water. Thin Starch Paste. A very dilute Solution of Caustic Alkali or Alkaline Carbonate, such as common Washing Soda. Solution of Hyposulphite of Sodium.

LEAD, SOLUBLE SALTS OF.

Sulphate of Sodium, Sulphate of Magnesium, Albumen, Emetics, and Purgatives, followed by Opium.

LIME, CHLORIDE OF.

Emetics, Albumen, Milk, Flour, Olive Oil.

MERCURY, SOLUBLE SALTS OF.

Albumen, Flour or Milk, Emetics. Four grains of Corrosive Sublimate require the White of *one* Egg only, as the Albumen must not be in excess.

OPIUM AND MORPHINE.

Emetics, Stomach Pump, strong Tea or Coffee, External Stimulants, Cold Affusion, Electro-Magnetism, *Compelled Exertion,* Artificial Respiration, and Sulphate of Atropine $\frac{1}{60}$ of a grain (subcutaneously), to be repeated if necessary.

PARAFFIN.

As for Carbolic Acid.

PHOSPHORUS.

Magnesia with Mucilage and Aromatic Spirit of Ammonia, followed by Oil of Turpentine.

NITRATE OF SILVER.

Common Salt (Chloride of Sodium), Albumen, Milk, Emetics.

STRYCHNINE.

See Alkaloids.

SULPHURETTED HYDROGEN OR THE ALKALINE SULPHIDES.

Dilute Solution of Chlorinated Soda or Lime (Bleaching Liquids), followed by the treatment of Ordinary Irritant Poisoning.

TABLE

Shewing the Proportion of active Ingredients contained in some important Preparations.

ACONITINE.

Unguentum Aconitinæ contains about gr. 8 of Aconitine in 1 ounce.

ANTIMONY.

Vinum Antimoniale contains gr. 1 of Tartar Emetic in fl. dr. iv.

ARSENIC.

Liquor Arsenii et Hydrargyri Iodidi (Donovan's) contains 1 per cent. of Iodide of Arsenium and 1 per cent. of Red Iodide of Mercury.

Liquor Arsenicalis contains 1 per cent. of Arsenious Acid.

Liquor Arsenici Hydrochloricus contains 1 per cent. of Arsenious Acid.

Liquor Sodii Arsenatis contains 1 per cent. of dried Arsenate of Sodium.

ATROPINE.

Unguentum Atropinæ contains gr. 8 of Atropine in oz. 1.

Liquor Atropinæ Sulphatis contains 1 per cent. of Sulphate of Atropine.

BELLADONNA.

Extractum Belladonnæ Alcoholicum contains 1 per cent. of the Alkaloids of Belladonna.

Extractum Belladonnæ Liquidum contains ·75 per cent. of the Alkaloids of Belladonna.

CANNABIS INDICA.

Tinctura Cannabis Indicæ contains gr. 1 of Extract in min. xxii.

COCAINE.

Unguentum Cocainæ contains 4 per cent. of Cocaine.

Injectio Cocainæ Hypodermica contains gr. 1 of Hydrochloride of Cocaine in min. xi.

CODEINE.

Syrupus Codeinæ contains gr. $\frac{1}{4}$ of Codeine Phosphate in fl. dr. j.

DIGITALIS.

Tinctura Digitalis contains about gr. 1 of dried leaves in min. ix.

Infusum Digitalis contains gr. 1 of dried leaves in min. clx.

HOMATROPINE.

Lamellæ Homatropinæ each contain gr. $\frac{1}{100}$ of Homatropine Hydrobromide.

IODINE.

Liquor Iodi Fortis contains about gr. 44 of Iodine in fl. oz. j.

Pigmentum Iodi contains about gr. 25 of Iodine in fl. oz. j.

IODIDE OF IRON.

Syrupus Ferri Iodidi contains about gr. $5\frac{1}{2}$ of Iodide of Iron in fl. dr. j.

MERCURY.

Unguentum Hydrargyri contains gr. 1 of Metallic Mercury in gr. 2.

Hydrargyrum cum Cretâ contains gr. 1 of Metallic Mercury in gr. 3.

Pilula Hydrargyri contains gr. 1 of Metallic Mercury in gr. 3.

PERCHLORIDE OF MERCURY.

Liquor Hydrargyri Perchloridi contains gr. 1 of Perchloride in fl. oz. ij.

Gargarisma Hydrargyri Perchloridi contains gr. 1 in fl. oz. v.

SUBCHLORIDE OF MERCURY.

Pilula Hydrargyri Subchloridi Composita contains gr. 1 of Calomel in gr. 5.

NUX VOMICA.

Tinctura Nucis Vomicæ contains gr. 1 of Strychnine in addition to Brucine in fl. oz. j.

STRYCHNINE.

Liquor Strychninæ Hydrochloridi contains 1 per cent. of Strychnine.

Syrupus Ferri Phosphatis cum Quinina et Strychninâ contains gr. $\frac{1}{32}$ of Strychnine in fl. dr. j.

OPIUM.

Pilula Saponis Composita contains gr. 1 of Opium in gr. 5.

Pilula Plumbi Composita contains gr. 1 of Opium and gr. 6 Acetate of Lead in gr. 8.

Pulvis Ipecacuanhæ Compositus contains gr. 1 of Opium in gr. 10.
Pulvis Opii Compositus contains gr. 1 of Opium in gr. 10.
Tinctura Opii contains gr. 1 of Opium in min. xv.
Pulvis Kino Compositus contains gr. 1 of Opium in gr. 20.
Pilula Ipecacuanhæ cum Scillâ contains gr. 1 of Opium in gr. 22.
Pulvis Cretæ Aromaticus cum Opio contains gr. 1 of Opium in gr. 40.
Tinctura Opii Ammoniata contains gr. 1 of Opium in min. xcvj.
Tinctura Camphoræ Composita contains gr. 1 of Opium in fl. oz. ss.
Linimentum Opii contains fl. oz. j. of Tincture in fl. oz. ij.

EXTRACT OF OPIUM.

Extractum Opii, gr. 2=about gr. 4 of Powdered Opium.
Extractum Opii Liquidum contains gr. 1 of Extract in min. xxv.

ACETATE OF MORPHINE.

Liquor Morphinæ Acetatis contains 1 per cent. of Acetate of Morphine.

Injectio Morphinæ Hypodermica fortior contains gr. 1 of Acetate of Morphine in min. vj.

Injectio Morphinæ Hypodermica (B.P.) contains gr. 1 of Tartrate of Morphine in min. xxii.

HYDROCHLORIDE OF MORPHINE.

Liquor Morphinæ Hydrochloridi contains 1 per cent. of Hydrochloride of Morphine.

Suppositoria Morphinæ contain gr. $\frac{1}{4}$ of Hydrochloride of Morphine in each Suppository.

Trochisci Morphinæ contain gr. $\frac{1}{36}$ of Hydrochloride of Morphine in each Lozenge.

Trochisci Morphinæ et Ipecacuanhæ contain gr. $\frac{1}{36}$ of Hydrochloride of Morphine and gr. $\frac{1}{12}$ of Ipecacuanha in each Lozenge.

Liquor Morphinæ Tartratis contains 1 per cent. of Tartrate of Morphia.

Tinctura Chloroformi et Morphinæ Composita contains min. $\frac{3}{4}$ of Chloroform, min. $\frac{1}{2}$ of Diluted Hydrocyanic Acid, and gr. $\frac{1}{11}$ of Morphine Hydrochloride in min. x.

PHOSPHORUS.

Pilula Phosphori contains about gr. 1 of Phosphorus in gr. 50.

Oleum Phosphoratum contains about gr. 1 of Phosphorus in min. c.

PREPARATIONS NOT INCLUDED IN THE BRITISH PHARMACOPŒIA.

DECOCTUM QUERCÛS.

℞ Corticis Quercûs.. .. .. .. oz. $1\frac{1}{4}$.
Aquæ destillatæ .. .. O j.

Boil for ten minutes, strain, and add sufficient distilled water to make 1 pint (see B.P., 1885).

Used in the preparation of Injectio Quercûs Composita.

DECOCTUM CINCHONÆ.

℞ Corticis Cinchonæ Rubris Contusæ.: oz. $1\frac{1}{4}$.
Aquæ destillatæ .. .. O j.

Boil for ten minutes, strain when cold, and add sufficient distilled water to make 1 pint (see B.P., 1885).

Used in the preparation of Haustus Cinchonæ Acidus and Haustus Cinchonæ cum Ammoniâ.

INFUSUM GENTIANÆ.

℞ Radicis Gentianæ Contusæ qr. ss. 120
Spiritûs Chloroformi .. .. .. min. c.
Aquæ bullientis. Oj

Macerate for 1 hour and strain.

Used in the preparation of Haustus Gentianæ Acidus, Haustus Gentianæ Alkalinus, and Haustus Potassii Iodidi Alkalinus.

TABLE

For Preparing Effervescing Draughts.

	Acidi Citrici.	Acidi Tartarici.	Succi Limonis.
Ammonii Carbonatis, gr. 20 = ...	gr. 24	25	fl. dr. vj.
Magnesii „ „ = ...	gr. 29	31	fl. dr. vij$\frac{1}{4}$.
Potassii „ „ = ...	gr. 17	18	fl. dr. iv$\frac{1}{4}$.
„ Bicarbonatis „ = ...	gr. 14	15	fl. dr. iij$\frac{1}{2}$.
Sodii „ „ = ...	gr. 17	18	fl. dr. iv$\frac{1}{4}$.

POSOLOGICAL TABLE.

Ages.	Proportional Quantities.	Doses.
For an Adult	Suppose the Dose to be One	Or 60 grs.
Under 1 year	Will require only $\frac{1}{12}$	„ 5 „
„ 2 years	„ „ „ $\frac{1}{8}$	„ $7\frac{1}{2}$ „
„ 3 „	„ „ „ $\frac{1}{6}$	„ 10 „
„ 4 „	„ „ „ $\frac{1}{4}$	„ 15 „
„ 7 „	„ „ „ $\frac{1}{3}$	„ 20 „
„ 14 „	„ „ „ $\frac{1}{2}$	„ 30 „
„ 20 „	The full Dose	„ 60 „

TABLE OF DOSES.

Acidi Acetici diluti .. .. fl. dr. j. to fl. dr. ij.
„ Benzoici .. .. .. gr. 5 to gr. 15.
„ Carbolici .. .. .. gr. 1 to gr. 3.
„ Gallici .. .. gr. 5 to gr. 15, or more.

Acidi Hydrobromici diluti ..	min. xv. to min. lx.
„ Hydrochlorici diluti ..	min. v. to min. xx.
„ Hydrocyanici diluti ..	min. ij. to min. vi.
„ Nitrici diluti	min. v. to min. xx.
„ Nitro-Hydrochlorici diluti	min. v. to min. xx.
„ Phosphorici diluti ..	min. v. to min. xx.
„ Salicylici	gr. 5 to gr. 20.
„ Sulphurici Aromatici..	min. v. to min. xx.
„ „ diluti ..	min. v. to min. xx.
„ Sulphurosi	fl. dr. ss. to fl. dr. j.
„ Tannici	gr. 2 to gr. 5.
Aloës Barbadensis	gr. 2 to gr. 5.
„ Socotrinæ	gr. 2 to gr. 5.
Aluminis	gr. 5 to gr. 10.
Ammonii Benzoätis	gr. 5 to gr. 15.
„ Bromidi	gr. 5 to gr. 30.
„ Carbonatis	gr. 3 to gr. 10.
„ Chloridi	gr. 5 to gr. 20.
Amyl Nitritis, by inhalation ..	min. ij. to min. v.
„ „ in mixtures	min. ss. to min. j.
Antimonii Tartarati—	
As a Diaphoretic and Expectorant	gr. $\frac{1}{24}$ to gr. $\frac{1}{8}$.
As an Emetic	gr. 1 to gr. 2.

Antifebrin	gr. 1 to gr. 3.
Antipyrin	gr. 5 to gr. 20.
Apomorphinæ Hydrochloridi (Hypodermically)—	gr. $\frac{1}{20}$ to gr. $\frac{1}{10}$.
As an Emetic	gr. $\frac{1}{10}$ to gr. $\frac{1}{4}$.
Argenti Nitratis	gr. $\frac{1}{4}$ to gr. $\frac{1}{2}$.
„ Oxidi	gr. $\frac{1}{2}$ to gr. 2.
Atropinæ Sulphatis— (Hypodermically)	gr. $\frac{1}{120}$ to gr. $\frac{1}{60}$.
Bismuthi Carbonatis	gr. 5 to gr. 20.
„ Citratis..	gr. 2 to gr. 5.
„ et Ammonii Citratis	gr. 2 to gr. 5.
„ Oxidi	gr. 5 to gr. 20.
„ Subnitratis	gr. 5 to gr. 20.
Boracis	gr. 5 to gr. 20.
Butyl-Chloral Hydratis ..	gr. 5 to gr. 20.
Caffeinæ..	gr. 1 to gr. 5.
„ Citratis	gr. 2 to gr. 10.
Calcii Chloridi	gr. 5 to gr. 15.
„ Hypophosphitis.. ..	gr. 3 to gr. 10.
Calcis Sulphuratæ	gr. $\frac{1}{4}$ to gr. 1.
Camphoræ	gr. 2 to gr. 5.
Capsici Pulveris	gr. $\frac{1}{2}$ to gr. 1.
Cerii Oxalatis	gr. 2 to gr. 10.
Chloral Hydratis	gr. 5 to gr. 20.
Chloroformi	min. j. to min. v.

Cocainæ Hydrochloridi ..	gr. $\frac{1}{5}$ to gr. $\frac{1}{2}$.
As a local anæsthetic, by painting or in a spray	Solutions varying from 1 to 20 per cent.
For Ophthalmic operations	A 2 or 4 per cent. solution dropped into the eye, or the lamellæ (gr. $\frac{1}{50}$), may be used.
Syr. Codeinæ	fl. dr. ss. to fl. dr. ij.
Codeinæ Phosp.	gr. $\frac{1}{4}$ to gr. 2.
Codeinæ..	gr. $\frac{1}{4}$ to gr. 2.
Confectionis Piperis	gr. 60 to gr. 120.
„ Sennæ	gr. 60 to gr. 120.
„ Sulphuris ..	gr. 60 to gr. 120.
Copaibæ	fl. dr. ss. to fl. dr. j.
Creasoti	min. j. to min. v.
Cretæ Præparatæ	gr. 10 to gr. 60.
Croton-Chloral Hydratis (*vide* Butyl-Chloral Hydratis)	gr. 5 to gr. 20.
Cubebæ Pulveris	gr. 30 to gr. 60.
Cupri Sulphatis	gr. $\frac{1}{4}$ to gr. 2.
Cusso	oz. $\frac{1}{4}$ to oz. $\frac{1}{2}$.
Decocti Aloës Compositi ..	fl. oz. ss. to fl. oz. ij.
„ Granati Radicis ..	fl. oz. ij. to fl. oz. iv.
„ Hæmatoxyli	fl. oz. j. to fl. oz. ij.
Digitalis Foliorum Pulveris	gr. $\frac{1}{2}$ to gr. 2.

Elaterini	gr. $\frac{1}{40}$ to $\frac{1}{10}$.
Elaterii	gr. $\frac{1}{10}$ to gr. $\frac{1}{2}$.
Ergotæ	gr. 20 to gr. 60.
Extracti Ergotæ (Hypodermically)	gr. 2 to gr. 5.
Extracti Aloës Barbadensis	gr. 1 to gr. 4.
„ Belladonnæ Viride	gr. $\frac{1}{4}$ to gr. 1.
„ „ Alcoholici	gr. $\frac{1}{4}$ to gr. 1.
„ Cannabis Indicæ ..	gr. $\frac{1}{4}$ to gr. 1.
„ Cascaræ Sagradæ ..	gr. 2 to gr. 8.
„ „ „ Liq.	fl. dr. ss. to fl. dr. j.
„ Cimicifugæ Liquidi	min. v. to min. xxx.
„ Cinchonæ Liquidi ..	min. v. to min. xv.
„ Cocæ Liquidi ..	fl. dr. ss. to fl. dr. j.
„ Colchici	gr. $\frac{1}{2}$ to gr. 1.
„ Colocynthidis Compositi	gr. 2 to gr. 5.
„ Ergotæ Liquidi ..	min. x. to min. xxx.
„ Euonymi Sicci ..	gr. 1 to gr. 2.
„ Filicis Liquidi ..	min. xlv. to min. xc.
„ Gelsemii Alcoholici	gr. $\frac{1}{2}$ to gr. 2.
„ HamamelidisLiquidi	min. v. to min. xv.
„ Hydrasti Liquidi ..	min. v. to min. xv.
„ Hyoscyami Viride	gr. 2 to gr. 8.
„ Jaborandi Liquidi..	min. v. to min. xv.
„ Jalapæ	gr. 2 to gr. 8.
„ Krameriæ	gr. 5 to gr. 15.

Extracti Nucis Vomicæ ..	gr. $\frac{1}{4}$ to gr. 1.
„ Opii..	gr. $\frac{1}{4}$ to gr. 1.
„ „ Liquidi	min. v. to min. xxx.
„ Pareiræ Liquidi ..	fl. dr. ss. to fl. dr. ij.
„ Physostigmatis ..	gr. $\frac{1}{4}$ to gr. 1.
„ Stramonii	gr. $\frac{1}{4}$ to gr. 1.
„ Taraxaci..	gr. 5 to gr. 15.
„ „ Liquidi ..	min. xxx. to fl. dr. ij.
Fellis Bovini Purificati ..	gr. 5 to gr. 15.
Ferri Arsenatis	gr. $\frac{1}{16}$ to gr. $\frac{1}{4}$.
„ Carbonatis Saccharatæ	gr. 10 to gr. 30.
„ et Ammonii Citratis ..	gr. 5 to gr. 10.
„ et Quininæ „ ..	gr. 5 to gr. 10.
„ Phosphatis	gr. 5 to gr. 10.
„ Redacti	gr. 1 to gr. 5.
„ Sulphatis	gr. 1 to gr. 5.
„ „ Exsiccatæ ..	gr. $\frac{1}{2}$ to gr. 3.
„ Tartarati	gr. 5 to gr. 10.
Glycerini	fl. dr. j. to fl. dr. ij.
„ Acidi Carbolici ..	min. v. to min. x. = about gr. 1 to gr. 2.
„ „ Tannici ..	min. x. to fl. dr. j. = about gr. 2 to gr. 12.
Guaiaci Resinæ Pulveris ..	gr. 5 to gr. 15.

Guaranæ	gr. 10 to gr. 60.
Homatropinæ Hydrobromidi	gr. $\frac{1}{8\,\overline{0}}$ to gr. $\frac{1}{2\,\overline{0}}$.
Hydrargyri cum Cretâ ..	gr. 1 to gr. 5.
„ Iodidi Rubri ..	gr. $\frac{1}{3\,\overline{2}}$ to gr. $\frac{1}{1\,\overline{6}}$.
„ Perchloridi ..	gr. $\frac{1}{3\,\overline{2}}$ to gr. $\frac{1}{1\,\overline{6}}$.
„ Subchloridi ..	gr. $\frac{1}{2}$ to gr. 5.
Hyoscyaminæ Sulphatis (Hypodermic Injection).. ..	gr. $\frac{1}{1\,2\,\overline{0}}$ to gr. $\frac{1}{4\,\overline{0}}$.
Infusi Aurantii	fl. oz. $\frac{1}{2}$ to fl. oz. j.
„ „ Compositi ..	fl. oz. $\frac{1}{2}$ to fl. oz. j.
„ Buchu	fl. oz. j. to fl. oz. ij.
„ Calumbæ	fl. oz. $\frac{1}{2}$ to fl. oz. j.
„ Caryophylli	fl. oz. j. to fl. oz. ij.
„ Cascarillæ	fl. oz. $\frac{1}{2}$ to fl. oz. j.
„ Chiratæ	fl. oz. $\frac{1}{2}$ to fl. oz. j.
„ Cinchonæ Acidi ..	fl. oz. $\frac{1}{2}$ to fl. oz. j.
„ Cuspariæ	fl. oz. j. to fl. oz. ij.
„ Digitalis	fl. dr. ij. to fl. dr. iv.
„ Ergotæ	fl. oz. j. to fl. oz. ij.
„ Gentianæ Simplicis ..	fl. oz. j. to fl. oz. ij.
„ „ Compositi	fl. oz. $\frac{1}{2}$ to fl. oz. j.
„ Krameriæ	fl. oz. $\frac{1}{2}$ to fl. oz. j.
„ Lupuli	fl. oz. j. to fl. oz. ij.
„ Quassiæ	fl. oz. $\frac{1}{2}$ to fl. oz. j.
„ Rhei	fl. oz. $\frac{1}{2}$ to fl. oz. j.
„ Rosæ Acidi	fl. oz. $\frac{1}{2}$ to fl. oz. j.
„ Senegæ	fl. oz. $\frac{1}{2}$ to fl. oz. j.

Infusi Sennæ	fl. oz. j. to fl. oz. ij.
„ Serpentariæ	fl. oz. ½ to fl. oz. j.
„ Uvæ Ursi	fl. oz. ½ to fl. oz. j.
Injectionis Apomorphinæ Hypodermicæ (contains gr. j. in min. 110)	min. v. to min. x.
Injectionis Ergotæ Hypodermicæ (contains about gr. j. in min. iij.)	min. iij. to min. x.
Injectionis Morphinæ Hypodermicæ B.P. (contains gr. j. in min. xxii.) . ..	min. ij. to min. v.
Injectionis Morphinæ Hypodermicæ Fort (contains gr. j. in min. vj.)	*min. ss. to min. iij.
Injectionis Morphinæ et Atropinæ Hypodermicæ (contains gr. $\frac{3}{10}$ of Tartrate of Morphine and gr. $\frac{1}{100}$ of Sulphate of Atropine in min. vj.)	*min. ij. to min. viij.
Injectionis Cocainæ Hypodermicæ (contains gr. x. in min. cx.)	min. ij. to min. v.
Iodi	gr. ½ gradually increased.
Iodoformi	gr. ½ to gr. 3.
Ipecacuanhæ—	
As an Expectorant ..	gr. ¼ to gr. 2.
As an Emetic	gr. 15 to gr. 30.

Jalapæ Pulveris	gr. 5 to gr. 20.
„ Resinæ	gr. 2 to gr. 5.
Kino	gr. 5 to gr. 20.
Liquoris Ammonii Acetatis	fl. dr. ij. to fl. dr. vj.
„ „ Citratis ..	fl. dr. ij. to fl. dr. vj.
„ Arsenicalis	min. ij. to min. viij.
„ Arsenici Hydrochlorici	min. ij. to min. viij.
„ Arsenii et Hydrargyri Iodidi (Donovan's Solution)	min. v. to min. xx.
„ Atropinæ Sulphatis	min. ½ to min. j.
„ Bismuthi et Ammonii Citratis.. ..	fl. dr. ss. to fl. dr. j.
„ Calcis	fl. oz. j. to fl. oz. iv.
„ „ Saccharati ..	min. xx. to fl. dr. j.
„ Ferri Acetatis ..	min. v. to min. xv.
„ „ Perchloridi ..	min. v. to min. xv.
„ „ Pernitratis ..	min. v. to min. xv.
„ Hydrargyri Perchloridi	fl. dr. ss. to fl. dr. j.
„ Magnesii Carbonatis	fl. oz. j. to fl. oz. ij.
„ Morphinæ Acetatis	min. x. to fl. dr. j.
„ „ Hydrochloridi	min. x. to fl. dr. j.
„ Morphinæ Tartratis	min. x. to min. lx.
„ Opii Sedativi ..	min. v. to min. xxx.

Liquoris Potassæ	min. x. to min. xxx.
„ Potassii Permanganatis	fl. dr. ij. to fl. dr. iv.
„ Sodæ Chlorinatæ ..	min. x. to min. xx.
„ Sodii Arsenatis ..	min. ij. to min. viij.
„ Strychninæ Hydrochloridi	min. ij. to min. viij.
Lithii Carbonatis	gr. 2 to gr. 5.
„ Citratis	gr. 5 to gr. 10.
„ „ Effervescentis..	fl. dr. j. to fl. dr. ij.
Magnesiæ Levis	gr. 10 to gr. 60.
„ Ponderosæ	gr. 10 to gr. 60.
Magnesii Carbonatis Levis ..	gr. 10 to gr. 60.
„ „ Ponderosæ	gr. 10 to gr. 60.
„ Sulphatis	gr. 30 to oz. ½.
„ „ Effervescentis	oz. ¼ to oz. 1.
Menthol..	gr. ½ to gr. 2.
Misturæ Ammoniaci	fl. oz. ss. to fl. oz. j.
„ Creasoti..	fl. oz. ½ to fl. oz. j.
„ Cretæ	fl. oz. j. to fl. oz. ij.
„ Ferri Compositæ ..	fl. oz. ½ to fl. oz. j.
„ Guaiaci	fl. oz. ½ to fl. oz. j.
„ Sennæ Compositæ	fl. oz. j. to fl. oz. ij.
Morphinæ Acetatis	gr. ⅛ to gr. ½.
„ Hydrochloridi ..	gr. ⅛ to gr. ½.
„ Tartratis	gr. ⅛ to gr. ½.
Nitroglycerini	gr. $\frac{1}{100}$.

Nitroglycerini Liquoris (contains 1 per cent.) ..	min. $\frac{1}{2}$ to min. ij.
„ Tabellæ (contain $\frac{1}{100}$ gr.)	1 or 2 tablets.
Nucis Vomicæ	gr. 1 to gr. 4.
Olei Copaibæ..	min. v. to min. xx.
„ Crotonis..	min. $\frac{1}{3}$ to min. j.
„ Morrhuæ	fl. dr. j. to fl. dr. iv.
„ Olivæ	fl. dr. j. to fl. dr. viij.
„ Ricini	fl. dr. j. to fl. dr. viij.
„ Terebinthinæ—	
As an Anthelmintic..	fl. dr. iij. to fl. dr. iv.
As a Diuretic	min. v. to min. x.
Opii Pulveris..	gr. $\frac{1}{4}$ to gr. 2.
Oxymellis Scillæ	fl. ss. to fl. dr. j.
Paraldehyde	min. xxx. to fl. dr. ij.
Pepsin..	gr. 5 to gr. 10.
Phenacetin	gr. 5 to gr. 10.
Pilocarpinæ Nitratis—	
(By the mouth)	gr. $\frac{1}{20}$ to gr. $\frac{1}{2}$.
(Hypodermically).. ..	gr. $\frac{1}{10}$ to gr. $\frac{1}{3}$.
Pilulæ Ferri	gr. 5 to gr. 15.
„ Hydrargyri	gr. 4 to gr. 8.
„ Plumbi cum Opio ..	gr. 2 to gr. 4.
„ Saponis Compositæ ..	gr. 2 to gr. 4.
Plumbi Acetatis	gr. 1 to gr. 5.

Podophylli Resinæ	gr. $\frac{1}{4}$ to gr. 1.
Potassii Acetatis	gr. 10 to gr. 60.
„ Bicarbonatis	gr. 5 to gr. 30.
„ Bromidi	gr. 5 to gr. 30.
„ Carbonatis	gr. 5 to gr. 20.
„ Chloratis	gr. 5 to gr. 15.
„ Citratis	gr. 10 to gr. 40.
„ Iodidi	gr. 5 to gr. 20.
„ Nitratis	gr. 5 to gr. 20.
„ Permanganatis ..	gr. 1 to gr. 3.
„ Sulphatis	gr. 10 to gr. 40.
„ Tartratis	gr. 60 to oz. $\frac{1}{2}$.
„ „ Acidæ ..	gr. 20 to gr. 60.
Pulveris Antimonialis	gr. 3 to gr. 6.
„ Catechu Compositi	gr. 10 to gr. 40.
„ Cretæ Aromatici ..	gr. 10 to gr. 60.
„ „ „ cum Opio	gr. 10 to gr. 40.
„ Elaterini Compositi (contains gr. 1 of Elaterine in gr. 40)	gr. 1 to gr. 4.
„ Glycyrrhizæ Compositi	gr. 60 to gr. 120.
„ Ipecacuanhæ Compositi	gr. 5 to gr. 15.
„ Jalapæ Compositi ..	gr. 20 to gr. 60.
„ Kino Compositi ..	gr. 5 to gr. 20.

Pulveris Rhei Compositi ..	gr. 20 to gr. 60.
„ Scammonii Compositi	gr. 10 to gr. 20.
Quininæ Hydrochloridi ..	gr. 1 to gr. 10.
„ Sulphatis, as a Tonic	gr. 1 to gr. 10.
„ „ as an Antipyretic	gr. 10 to gr. 30.
Salicini	gr. 3 to gr. 20.
Santonini	gr. 2 to gr. 5.
Scammoniæ	gr. 5 to gr. 10.
Scammonii Resinæ	gr. 3 to gr. 8.
Scillæ Pulveris	gr. 1 to gr. 3.
Sinapis, as an Emetic	From a dessertspoonful to a tablespoonful.
Sodæ Tartaratæ	oz. $\frac{1}{4}$ to oz. $\frac{1}{2}$.
Sodii Arsenatis	gr. $\frac{1}{40}$ to gr. $\frac{1}{10}$.
„ Bicarbonatis	gr. 5 to gr. 30.
„ Bromidi	gr. 5 to gr. 30.
„ Carbonatis	gr. 5 to gr. 30.
„ Hypophosphitis	gr. 3 to gr. 10.
„ Iodidi	gr. 5 to gr. 20.
„ Phosphatis	gr. 30 to gr. 120.
„ „ Effervescentis	oz. $\frac{1}{4}$ to oz. $\frac{1}{2}$.
„ Salicylatis	gr. 10 to gr. 30.
„ Sulphatis Effervescentis	oz. $\frac{1}{4}$ to oz. $\frac{1}{2}$.

Sodii Sulphocarbolatis	gr. 5 to gr. 15.
Spiritûs Ætheris	min. xx. to min. xl.
,, ,, Nitrosi ..	min. xx. to min. xl.
,, Ammoniæ Aromatici	min. xx. to min. xl.
,, ,, Fœtidæ..	min. xx. to min. xl.
,, Armoraciæ Compositi	fl. dr. j. to fl. dr. ij.
,, Camphoræ	min. v. to min. xx.
,, Chloroformi	min. v. to min. xx.
Strychninæ	gr. $\frac{1}{60}$ to gr. $\frac{1}{15}$.
Succi Belladonnæ..	min. v. to min. xv.
,, Conii	fl. dr. ss. to fl. dr. ij.
,, Hyoscyami	fl. dr. ss. to fl. dr. j.
,, Scoparii	fl. dr. j. to fl. dr. ij.
,, Taraxaci	fl. dr. j. to fl. dr. ij.
Sulphonal	gr. 10 to gr. 30.
Sulphuris Sublimati	gr. 20 to gr. 60.
Syrupi Chloral (contains gr. 10 in fl. dr. j.) ..	fl. dr. ss. to fl. dr. ij.
,, Ferri Iodidi	min. xx. to fl. dr. j.
,, ,, Phosphatis ..	min. x. to fl. dr. j.
,, ,, ,, Compositi	min. x. to fl. dr. j.
,, ,, ,, cum Quinina et Strychnina	fl. dr. ss. to fl. dr. j.
,, Rhei..	fl. dr. ss. to fl. dr. ij.
,, Scillæ	fl. dr. ss. to fl. dr. j.
,, Sennæ	fl. dr. j. to fl. dr. ij.

Terebene	.. ..	min. v. to min. xv.
Terebinthinæ Chiæ	.. ..	gr. 5 to gr. 10.
Theinæ	.. ..	gr. 1 to gr. 5.
Tincturæ Aconiti	.. .. ..	min. v. to min. xv.
„ Aloës	.. .. ..	fl. dr. ss. to fl. dr. j.
„ Asafœtidæ	.. ..	fl. dr. ss. to fl. dr. j.
„ Belladonnæ	.. ..	min. v. to min. xv.
„ Benzoini Compositæ	.. .. ..	fl. dr. ss. to fl. dr. j.
„ Buchu	.. .. ..	fl. dr. ss. to fl. dr. j.
„ Calumbæ	.. ..	fl. dr. ss. to fl. dr. j.
„ Camphoræ Compositæ	.. .. ..	fl. dr. ss. to fl. dr. j.
„ Cannabis Indicæ	..	min. v. to min. xv.
„ Cantharidis	.. ..	min. v. to min. xv.
„ Capsici	.. .. ..	min. v. to min. xv.
„ Catechu	.. .. ..	fl. dr. ss. to fl. dr. j.
„ Chiratæ	.. .. ..	fl. dr. ss. to fl. dr. j.
„ Chloroformi et Morphinæ Compositæ		min. v. to min. xv.
„ Cimicifugæ	.. ..	min. xxx. to fl. dr. j.
„ Cinchonæ	.. ..	fl. dr. ss. to fl. dr. j.
„ „ Compositæ		fl. dr. ss. to fl. dr. j.
„ Colchici Seminum		min. v. to min. xv.
„ Conii	.. .. ..	min. xxx. to fl. dr. j.
„ Cubebæ	.. .. ..	fl. dr. ss. to fl. dr. j.
„ Digitalis	.. ..	min. v. to min. xv.

Tincturæ	Ergotæ Ammoniatæ	fl. dr. ss. to fl. dr. j.
,,	Ferri Perchloridi..	min. v. to min. xv.
,,	Gelsemii	min. v. to min. xv.
,,	Gentianæ Compositæ	fl. dr. ss. to fl. dr. j.
,,	Guaiaci Ammoniatæ	fl. dr. ss. to fl. dr. j.
,,	Hamamelidis ..	fl. dr. ss. to fl. dr. j.
,,	Hydrastis	fl. dr. ss. to fl. dr. j.
,,	Hyoscyami	fl. dr. ss. to fl. dr. j.
,,	Iodi	min. ij. to min. v.
,,	Jaborandi	fl. dr. ss. to fl. dr. j.
,,	Jalapæ	fl. dr. ss. to fl. dr. j.
,,	Kino	fl. dr. ss. to fl. dr. j.
,,	Krameriæ	fl. dr. ss. to fl. dr. j.
,,	Lobeliæ Æthereæ..	min. v. to min. xv.
,,	Lupuli	fl. dr. ss. to fl. dr. j.
,,	Nucis Vomicæ ..	min. v. to min. xv.
,,	Opii	min. v. to min. xv. min. xx. to min. xxx.
,,	,, Ammoniatæ ..	fl. dr. ss. to fl. dr. j.
,,	Podophylli	min. v. to min. xv.
,,	Quininæ	fl. dr. ss. to fl. dr. j.
,,	,, Ammoniatæ	fl. dr. ss. to fl. dr. j.
,,	Rhei Compositæ—	
	As a Stomachic..	fl. dr. ss. to fl. dr. j.
	As a Purgative..	fl. dr. ij. to fl. dr. iv.
,,	Scillæ	min. v. to min. xv.

Tincturæ Senegæ	fl. dr. ss. to fl. dr. j.
„ Sennæ Compositæ	fl. dr. ss. to fl. dr. j. fl. dr. ij. to fl. dr. iv.
„ Serpentariæ	fl. dr. ss. to fl. dr. j.
„ Stramonii	min. v. to min. xv.
„ Strophanthi	min. v. to min. xv.
„ Valerianæ Ammoniatæ	fl. dr. ss. to fl. dr. j.
„ Zingiberis	fl. dr. ss. to fl. dr. j.
Vini Antimonialis ..	min. x. to min. xxx. fl. dr. ij. to fl. dr. iv.
„ Colchici	min. x. to min. xxx.
„ Ferri	fl. dr. j. to fl. dr. iv.
„ „ Citratis	fl. dr. j. to fl. dr. iv.
„ Ipecacuanhæ—	
As an Expectorant ..	min. x. to min. xxx.
As an Emetic	fl. dr. iv. to fl. dr. vj.
„ Quininæ	fl. oz. ss. to fl. oz. j.
Zinci Acetatis—	
As a Tonic	gr. 1 to gr. 2.
As an Emetic	gr. 10 to gr. 20.
„ Chloridi	gr. ½ to gr. 2.
„ Oxidi	gr. 3 to gr. 10.
„ Sulphatis—	
As a Tonic	gr. 1 to gr. 3.
As an Emetic	gr. 10 to gr. 30.
„ Valerianatis	gr. 1 to gr. 3.

INSTRUCTIONS

FOR

CASE-TAKING

IN THE

MEDICAL WARDS.

INSTRUCTIONS FOR CASE-TAKING IN THE MEDICAL WARDS.

Name. Sex. Age. Residence. Occupation.

FAMILY HISTORY.

Parentage.—Age of parents. If living, state if they are healthy, or if subject to any and what disease. If dead, state causes of death, and the age they attained.

Brothers and Sisters.—Number. Ages. Diseases. Causes of Death.

Hereditary Taint.—Enquire especially for evidence of following classes of disease in members of family :—phthisis, rheumatism, gout, syphilis, cancer, nervous disorders (*e.g.*, epilepsy, mania, etc.).

PREVIOUS HISTORY.

Enumerate previous illnesses, with times of their occurrence. Habitual state of health.

Habits and Mode of Life.—Abstemious or otherwise. Variety of habitual food and drink. Occupation—sedentary or active, how long followed, with or without interruption. Residence, where, and for how long. Climate, damp or dry.

Civil State.—Date of marriage. Number, ages, and diseases of children.

N.B.—If a woman, enquire as to date of first appearance and regularity of catamenia, number and date of pregnancies, miscarriages (if any).

PRESENT ILLNESS.

Date and mode of onset; sudden or gradual; and to what cause attributed.

State how long work has been left off, and how long patient has taken to bed.

Enquire as to symptoms, giving them in the order of their occurrence, and their duration. Enquire especially as to rigors, pains (their precise seat and character), appetite, state of bowels and urine, cough,

expectoration, loss of flesh, sweating, muscular weakness, sleep, disorders of special senses, etc.

PRESENT CONDITION.

Record rate of pulse and respirations, and temperature.

Bodily conformation. Stature. Complexion. Temperament.

N.B.—Note any deformity (rickets, spinal curvature, etc.).

Position in bed.

Expression of Face.—Pupils dilated or contracted, equal or unequal; arcus senilis; œdema of conjunctivæ; strabismus.

Colour and condition (dry or moist) of skin and visible mucous membranes, lips, conjunctivæ, finger-nails—whether pale, congested, livid, or jaundiced. Cutaneous eruptions—their character, seat, time of appearance, and duration.

Presence or absence of œdema, noting where it is marked.

Voice. Cough. Expectoration.

SUBJECTIVE SYMPTOMS.

State these as nearly as possible in patient's own words.

Morbid sensations—their seat and character.

Pain.—Time of occurrence—paroxysmal or continuous—character (dull, aching, shooting, etc.),—precise seat,—whether influenced by position or movement.

Numbness. Tingling. Formication.

Diplopia. Dimness of Sight. Disorders of hearing, smelling, etc.

Vertigo. Palpitation. Breathlessness (whether on exertion or without). Fainting.

Sense of heat and cold.

Hunger. Thirst. Nausea. Dysphagia.

OBJECTIVE SYMPTOMS.

Circulatory System.

Pulse.—Its rate, volume, regularity, and character [compare pulse at both wrists]; visibility of pulsation and

tortuosity of vessel at wrists, temples, and in supraclavicular regions.

Venous turgescence—pulsation.

HEART.

Inspection.—Position of apex-beat; character and extent of impulse (heaving, undulatory, diffused, etc.); Epigastric pulsation.

Palpation.—Confirm the foregoing. Cardiac thrill (seat of). Friction fremitus.

Percussion.—Area of Cardiac dulness.

Auscultation. — Sounds — character, rhythm, accentuation, doubling, muffling, etc. Murmurs—their relation to the sounds; where best heard, where conducted. Whether audible in carotids, axilla, and back.

Friction — extent, character, and rhythm.

RESPIRATORY SYSTEM.

LARYNX.

Laryngoscopic appearances.

CHEST.

Inspection.—Form, symmetry, prominence of superficial veins, expansion and ele-

vation—note any difference in movement of the two sides. Laboured breathing (dyspnœa, orthopnea). Note any retraction of soft parts in inspiration.

Measurement.—Cyrtometer.

Palpation.—Confirm facts ascertained by inspection. Fremitus—vocal, respiratory, rhonchal, frictional.

Percussion.—Resonance, area, and character; dulness, absolute or relative; mark out *precise* limits, and note whether influenced by position—"Cracked-pot sound."

Auscultation.—Vocal resonance; bronchophony, pectoriloquy, ægophony. Character of respiratory murmur; comparative duration and intensity of inspiration and expiration. [Breath sounds may be vesicular, weak, harsh, bronchial, tubular, amphoric, cavernous.]

"Bell-sound" of pneumothorax.

Adventitious Sounds.—Note particularly the regions where they are audible.

(*a*) *Moist Sounds or Râles.*—Crepitant, sub-crepitant, crackling (large or small); gurgling. Whether audible with inspiration or expiration, or both; whether modified by coughing or not.

(*b*) *Dry Sounds or Rhonchi.*—Sonorous, sibilant.

(*c*) *Friction Sounds.*

(*d*) Metallic tinkling.

Succussion.—Splashing sound.

[*Note.*—Remark any aneurysmal bruit that may be audible in any part of the chest, describing its precise seat and extent.]

DIGESTIVE SYSTEM.

Tongue.—Shape, size, colour; dry or moist; character and situation of fur. Appearance of papillæ.

Fauces and Tonsils.—Condition of.

Deglutition.—Difficulty in.

Examination of Abdomen.—State of parietes; obesity. Lineæ atrophicæ. Distension or retraction. Note precise region of any swelling. Prominence of superficial veins, and their direction. Condition of umbilicus. Measurements.

Palpation.—Tenderness: its exact seat; deep, superficial, local, general. Fluctuation. Gurgling, resistance, abdominal pulsation, tumours. Tâche—its characters.

Percussion.—Tympanites. Dulness—its exact region, and whether altered by change of position. [Hypogastric dulness, sometimes due to distended bladder.]

Auscultation.—Aortic and aneurysmal bruit. Pulsation of fœtal heart; uterine bruit.

Stomach.—Character of Vomit. (See Appendix.) Distension.—Area of resonance, etc.

Intestines.—Character and frequency of stools.

Liver.—Area of dulness; outline and surface. Note any irregularity, or local and general enlargement. "Sense of displacement."

Spleen.—Area of dulness: whether felt below ribs.

NERVOUS SYSTEM.

See "Subjective Symptoms."

Paralysis.—Partial or general, muscles involved. If of long standing note presence of wasting, contractions, degree of electric irritability. Faradaism. Galvanism.

Anæsthesia.—Distribution of.

Hyperæsthesia.—Distribution of.
Altered sensibility to heat and cold.

Spasm.—Tonic, clonic, muscles involved.
Loss of memory, speech—alterations in.
Aphasia.
Reflex irritability—"tendon reflex."

Ophthalmoscopic Examination.

URINARY SYSTEM.

Incontinence. Retention. Frequency and difficulty of micturition. Pain.

For examination of Urine, see Appendix.

APPENDIX.

Examination of Urine, Sputum, and Vomited Matters.

URINE.

Quantity passed in 24 hours. Colour, odour, specific gravity, reaction. Deposits, their colour and character.

Chemical Examination.—Urea, urates, phosphates, chlorides, albumen, pus, blood, bile, mucus, sugar.

Microscopical Examination.—(Note time since urine has been passed), crystals, form and nature; amorphous deposits. Tube casts—their form and character. Epithelium; blood, pus, and mucus. Corpuscles. Spermatozoa. Bacteria. Foreign bodies.

[N.B.—Make sketches of microscopical appearances.]

Quantitative Estimation of urea; sugar.

SPUTUM.

Character; consistency; colour; odour; aeration. Microscopical characters—pus; blood; elastic tissue.

VOMITED MATTERS.

Reaction; appearance; contents; presence of sarcinæ, etc.

Quantity—when vomited.

Note.—For systematic record in the Hospital Case Book it is convenient to adopt the order of enquiry much as here given. But in "taking a case" it is recommended to examine first the present condition of the patient (p. 103 *et seq.*), and then to enquire into the previous history, etc., as points for such enquiry will necessarily be raised in the examination. It need hardly be pointed out that it is by no means necessary in *every* case to go seriatim and in detail through all the signs here indicated; but it must be borne in mind that often *negative* signs are as valuable in diagnosis as positive.

INSTRUCTIONS

FOR

CASE-TAKING

IN THE

SURGICAL WARDS.

INSTRUCTIONS FOR CASE-TAKING IN THE SURGICAL WARDS.

A well-recorded case should embrace *all* the facts bearing on the injury or disease under consideration, and *no others*.

Before subjecting a patient to a prolonged examination, it should be ascertained whether he is in a fit condition to answer questions, and to bear the necessary manipulations. He may be delirious, comatose, or so exhausted as to render any examination highly injurious.

When the patient is unable to answer questions, the friends should be interrogated.

Every case should have this heading—"Disease: Result;" it greatly facilitates subsequent reference.

It need not necessarily be filled in till after the completion of the case.

GENERAL HISTORY.

Name. Age. Sex.
Single, married, or widow.
Residence.
Occupation.
Date of Admission (day of month and year).
Ward and Bed.
Under whose care.
Any obvious lesion or external morbid appearance: *e.g.*, fractured skull and coma.

HISTORY OF PRESENT ILLNESS.

How long ill.
When left off work.
How long in bed.
How did it begin—subsequent symptoms in sequence.
What was the cause.
Treatment before admission.

In cases of Accident or Urgency.

Mode of occurrence and nature of accident or injury. Part injured. If a wound, its characters.

Its exact date: in urgent cases stating hour.

Was there consequent insensibility, pallor, coldness of surface, hæmorrhage, vomiting, etc.

Drunk or sober.

Previous treatment.

PREVIOUS HISTORY.

Usual state of health (loss of flesh, weight, colour, etc.).

Previous diseases (dates and duration; if necessary give symptoms), especially as to Phthisis, Syphilis, Hysteria, Malaria, Rheumatic Fever and Gout.

Temperate, or excesses and irregularities.

Food, drink, tobacco, clothes, etc.

Social condition and mode of life.

How long followed present occupation.

Residence at home, abroad, or in India.

Exposure to extremes of weather (heat, cold, wet, etc.).

FAMILY HISTORY.

What diseases are known to run in the family: especially as to Phthisis, Syphilis, Gout, and Cancer.

Diseases of parents and grandparents. Their respective ages and causes of death.

Diseases, etc., of brothers, sisters, and their children.

Matrimonial life (number of children, their diseases, causes of death, etc., miscarriages).

PRESENT STATE.

This part of the case should be taken as soon after admission as possible; and in all cases of *accident* or *urgency*, a short account of the patient's *condition on admission* should be written at once.

Such points as the following are of importance:—

Sensible or insensible.
Pale or florid.
Degree of illness, the look of pain, etc.
Dull, apathetic or excited.
Drunk or sober.
Vomiting or purging.
Character of vomit.
Paralysis (especially as to limbs and face).
Convulsions, spasms, tremors, etc.

Also the Pulse and Respirations (the number per minute, character). Temperature. Tongue (especially movements, colour, froth, lacerations, etc.). Odour of breath. Pupils (size, and action to light). State of Special Senses. Hæmorrhage. Escape of cerebro-spinal fluid, or blood, from the ear or nose. Sub-conjunctival ecchymosis.

In the event of a patient being brought to the Hospital drunk or senseless, a most important question arises, fraught with all

sorts of disagreeable possibilities, Is he simply drunk—or dying?

In taking the Present State, the first thing is to make a *General Inspection* of the patient, attention being directed also to any obvious special symptoms, such as dyspnœa, cough, vomiting, etc.

Aspect.—Healthy or otherwise. Sensible or insensible. Mental State. Nutrition and conformation. Manner. Paralysis, tremors or rigors. Delirious or prostrated. Confined to bed or about as usual. Decubitus, attitude or gait. Orthopnœa. Position and movements of limbs, etc.

Countenance.—Anæmic, plethoric, cachetic, jaundiced, hectic, cyanotic, pasty, œdema of eyelids, paralysis (partial or general), tic, sordes (lips and gums).

Expression. — Cheerful, melancholic, apathetic, anxious, frightened, drawn as if in pain, 'facies hippocratica,' indicating approaching death.

Eyes.—Open or shut, fixed or oscillating, squint, pupils (size, action to light, etc.), blood vessels, arcus senilis, proptosis.

Skin.—Colour, hot or cold, dry or moist, eruptions, anasarca.

The next step comprises examination by *Interrogation.*

In eliciting the history of the case, it will have been ascertained whether the disease is *acute* or *chronic*—a most important consideration. It has now to be localized or recognized as one of the general diseases. For this purpose ask the following questions:—

(*a*) What do you complain of?
(*b*) Where do you feel pain?

By these questions, together with such subsidiary ones as naturally spring from them, the probable seat of the disease is determined.

A concise description of the lesion or diseased part should be given, together with such negative statements as seem likely to have a practical bearing.

Information as to the following points should be taken:—

Pulse.—Frequency. Rhythm:—Regular, irregular, or intermittent.

Volume:—Large, full, small, thready.

Force and tension:—Strong, weak, extinct, soft, hard, wiry, compressible, and incompressible.

Special characters:—Rigid, tortuous, bounding, hammering (Corrigan's), jerky, undulatory, thrilly, and dicrotic.

Respiration.—Frequency, easy or difficult (dyspnœa), orthopnœa, painful, laboured, short or hurried, deep or shallow, abdominal or thoracic.

Temperature.—As recorded by the thermometer, usually in the axilla (normal 98·4 F.).

Tongue.—Moist or dry, clean or furred, its colour, smooth, rough, or fissured, frothy, tremulous movements, ulcers, or lacerations.

Appetite.—Good, bad, or indifferent, anorexia, thirst, solids or fluids, relish of food, voracity, nausea, vomiting, etc.

Bowels.—Regularity, frequency, nature of motions, constipation, diarrhœa, flatus, and borborygmi. Fæces should be inspected in certain cases.

Urine.—Total amount in 24 hours (normal 30 to 50 oz.). General appearance and colour, sediments, reaction, sp. gr., albumen (casts), sugar, urates, phosphates, pus, mucus, blood, etc.

Menstruation.—Regularity, how often, how much, nature of discharge, pains.

TREATMENT.

Hygienic, dietetic and medicinal, stimulants, external applications. Prescriptions should be entered in the margin. Any apparatus or instrumentation should be clearly described.

In the event of an operation, note the successive steps, anæsthetic, incisions, instruments, hæmorrhage and means for arresting it, number and kind of ligatures, unexpected occurrences, dressing, antiseptic precautions, condition of patient on removal to bed and on recovery from anæsthesia, vomiting, etc.

When a diseased part has been removed, a description of the morbid appearances should be given.

PROGRESS.

The frequency of the requisite observations will depend on the nature of the case. In very urgent cases, daily or hourly reports should be made; but in chronic cases, one or two observations a week will suffice.

The pulse, respirations, and temperature should be taken every morning and evening in all *acute* cases, especially after an operation—a tabular record being kept.

In noting the progress from time to time the same general plan should be followed as in taking the Present State. Enquiry should be made respecting the pulse, respiration, temperature, tongue, appetite, bowels, pain, sleep, urine, and menstruation.

Any change in the patient's state or new phenomena are to be noted, and the effects of medicines enquired into; regard being paid to the accounts of the patient and the attendants, as well as to the phenomena actually observed by the medical man.

After an operation enquiries should be directed first to the patient's general condition, and secondly to the condition of the wound.

Under the former head look out for headache, malaise, nausea, vomiting, feeling of illness, chills, rigors, feverishness, etc. Such phenomena often give warning of morbid processes commencing in

the wound, but as yet invisible to the naked eye.

Under the latter, note the date of each change of dressing and any alterations made; the amount and character of the discharge; oozing or hæmorrhage; condition of edges of wound, their degree of apposition and surrounding inflammation; establishment of suppuration; the process of repair—first intention, granulations and their nature; removal of sutures, ligatures, and drainage tubes.

RESULT.

Its date and nature; patient's condition on leaving hospital—cured, relieved, or not benefited. Left for his home or convalescent institution. In the event of death, give a short account of the phenomena immediately preceding. State whether autopsy or not; if so, give a full account. Having witnessed the case from beginning to end, it is an excellent practice to re-write and condense the notes, with the aid of the new light thus brought to bear, and to add a short commentary on the most noteworthy features.

Mitchell and Hughes, Printers, 140 Wardour Street, W.

3 2/4 - 4 g

3.50 fat. 6.5 sugar 1.25 protein

THE MIDDLESEX HOSPITAL.—DIET TABLE, JANUARY, 1899.

DIETS.	DAILY.	BREAKFAST, 6 A.M.	LUNCHEON, 10.30 A.M.	DINNER, 12 NOON.	SUPPER, 7 P.M.	EXTRAS.
LIQUID	2 pints of Milk. 1 pint of Beef Tea.					Chops, ¼ lb. each when trimmed (Mutton or Pork). Steaks: Rump Steak ½ lb., without bone. Chicken, a quarter. Sausages, ½ lb. Rabbit, a quarter. Tripe, ½ lb. Bacon, 3 ozs. Soup, for Supper, 1¼ lbs. Meat, 16 ozs. of Vegetables, 4 ozs. Grain, to 5 diets. Seasoning to palate. Broth, without Meat, 1 lb. of lean Mutton, with bone, to a pint. Chicken Broth, ½ Chicken to a pint. Oysters. Greens. Suet Pudding:—1 oz. of Beef Suet. 2 ozs. of Flour. Custard Pudding:—½ oz. of Sugar. 1 pint of Milk. 2 Eggs. Arrowroot. Sago. Oatmeal Gruel, 3 ozs. to a pint. Eggs. Jellies. Oranges. Lemons.
MILK	10 ozs. Bread.	½ pint of Milk.	½ pint of Beef Tea.	ALTERNATE DAYS. Pudding. Egg. Milk. Sugar. 1½ oz. Rice. 1 1 pint. ½ oz. 1½ oz. Tapioca ½ 1 pint. ½ oz. or Sago.	½ pint of Milk. 1 pint of Soup.	
FISH	10 ozs. Bread.	½ pint of Milk.	½ pint of Beef Tea.	8 ozs. of Fish (Whiting, Haddock, Cod, Plaice, or Brill), Sole if specially ordered. ½ lb. of Potatoes.	½ pint of Milk. 1 pint of Soup.	
ORDINARY	10 ozs. Bread.	½ pint of Milk.	½ pint of Beef Tea.	Men, 6 ozs., Women, 3 ozs., Cooked Meat, without bone; 3 days a week roast, 3 days a week boiled, 1 day a week stew, with Vegetables. ½ lb. of Potatoes.	½ pint of Milk. 1 pint of Soup.	
MUTTON BROTH ...	10 ozs. Bread.	½ pint of Milk.	½ pint of Beef Tea.	8 ozs. Neck of Mutton, served in 1 pint of Broth, with Vegetables, and 2 ozs. of Barley. ½ lb. of Potatoes.	½ pint of Milk. 1 pint of Soup.	
CONVALESCENT ...	10 ozs. Bread.	½ pint of Milk.	½ pint of Beef Tea.	Men, 6 ozs., Women, 4 ozs., Cooked Meat, without bone, on week days; and of Beef, roast and boiled, alternately on Sundays. ½ lb. of Potatoes.	½ pint of Milk. 1 pint of Soup.	

STIMULANTS.

Wine. Porter.
Spirits. Bottled Stout, ½ pints.
Draught Ale. Ditto, pints.
Bottled Ale.

CHILDREN	DAILY ...	Bread, 6 ozs. Milk, 1 pint. Pudding (Rice or Sago), 2 ozs.	Cane Sugar, 1 oz. Jam, or Stewed Fruit, 1 oz. Butter, ½ oz.	EXTRAS	Mutton or Veal Broth (1 lb. to 1 pint). Fish, 3 ozs. Potatoes. Minced Meat. Rabbit or Chicken (¼ of a Chicken or ¼ of a Rabbit to serve for 2 diets). Greens. Eggs.

NOTES.

A.—Every Patient admitted into the Hospital is placed upon Milk Diet, until a Diet is otherwise ordered.
B.—Diets ordered by the Physicians and Surgeons are to be continued until changed by subsequent orders.
C.—The Diet Cards shall be submitted to the Physicians or Surgeons for revision, and are required to be initialled by them once a week.
D.—Saturday Diets are continued until Monday, and cannot be altered except in cases of emergency.
E.—The Sisters shall use their discretion in drawing Pudding and Potatoes for the Patients in their respective Wards, while not exceeding the allowance laid down in the Diet Table.
F.—In all cases where the Diet of a Patient has for some reason to be omitted, the Sister should send notice to the Clerk's Office as soon as possible, and in the event of such a Diet being subsequently required, she may requisition it later in the day. (Forms will be provided for this purpose.)
G.—No Extras or Beer can be issued unless the Physician or Surgeon initials the order on the "Extra Diets Sheet" at least once a week.
H.—The Dispenser is not to issue Stimulants unless the Physician or Surgeon initials the order on the "Stimulant Sheet" at least once a week.
I.—A Return of Stimulants supplied to be sent in weekly to the Medical Committee by the Dispenser.
J.—The above Dietary does not necessarily apply to the Patients in the Caner Wards, who are allowed special privileges according to the exigencies of the cases.

PHARMACOPŒIA

OF THE

MIDDLESEX HOSPITAL